arXiv:0806.0606v4 [math.DG] 30 Jul 2009

LARGE COMPLEX STRUCTURE LIMITS, QUANTIZATION AND COMPACT TROPICAL AMOEBAS ON TORIC VARIETIES

THOMAS BAIER, CARLOS FLORENTINO, JOSÉ M. MOURÃO
AND JOÃO P. NUNES

Abstract. We consider toric deformations of complex structures, described by the symplectic potentials of Abreu and Guillemin, with degenerate limits of the holomorphic polarization corresponding to the toric Lagrangian fibration, in the sense of geometric quantization. This allows us to interpolate continuously between quantizations in the holomorphic and real polarizations and show that the monomial holomorphic sections of the prequantum bundle converge to Dirac delta distributions supported on Bohr-Sommerfeld fibers.

Under these deformations, the original toric metric is deformed to non-equivalent toric metrics. We use these families of toric metric degenerations to study the limit of compact hypersurface amoebas and show that in Legendre transformed variables they are described by tropical amoebas. For a particularly relevant one-parameter family of deformations, the limiting tropical amoebas are compact and live naturally in the polytope, image of the moment map. We believe that our approach gives a different, complementary, perspective on the relation between complex algebraic geometry and tropical geometry.

Contents

1. Introduction and main results 1
1.1. Geometric quantization of toric varieties 2
1.2. Compact tropical amoebas 5
2. Preliminaries and notation 6
3. Quantization in a real polarization 9
4. Large toric complex structure limits of Kähler polarizations 12
5. Large complex structure limits of holomorphic sections and BS fibers 14
6. Compact tropical amoebas 17
6.1. Limit versus tropical amoebas 17
6.2. Implosion of polytopes versus explosion of fans 23
6.3. Amoebas associated to geometric quantization 24
6.4. Relation to other aspects of degeneration of complex structures 29
References 30

1. Introduction and main results

Studying an infinite-dimensional class of 1-parameter families of toric complex structures on a smooth toric variety, we investigate large complex structure limit points corresponding to holomorphic Lagrangian distributions degenerating to the real Lagrangian torus fibration defined by the moment map. We use ideas of geometric quantization rather than a purely algebro-geometric construction of a compactification of the moduli space of complex structures. This approach permits us to obtain the following main results. Let P be a Delzant polytope and let X_P be the associated compact toric variety [De]. Let $\psi \in C^\infty(P)$ be a smooth function with positive definite Hessian on P. Such ψ defines a compatible complex structure on X_P (see (2)). Then,

1. In Theorem 1.1, we determine the weakly covariantly constant sections of the natural line bundle on X_P with respect to the (singular) real polarization defined by the Lagrangian fibration given by the moment map μ_P. In particular, we see that they are naturally indexed by the integer points in the polytope P.

2. We show in Theorem 1.2 that the family of Kähler polarizations given by the deformations associated to $s\psi$ for $s \in \mathbb{R}^+$, converges as $s \to +\infty$ to the real polarization, independently of the direction of deformation.

3. Theorem 1.3 states that the holomorphic monomial sections of the natural line bundle converge, also for all ψ, to the Dirac delta distributions supported on the corresponding Bohr-Sommerfeld orbit of Theorem 1.1.

For the class of symplectic toric manifolds, this solves the important question, in the context of geometric quantization, on the explicit link between Kähler polarized Hilbert spaces and real polarized ones, in particular in a situation where the real polarization is singular. For a different, but related, recent result in this direction see [BGU].

4. We show that the compact amoebas [GKZ, FPT, Mi] of complex hypersurface varieties in X_P converge in the Hausdorff metric to tropical amoebas in the (ψ-dependent) variables defined from the symplectic ones via the Legendre transform

$$\begin{array}{rcl} \mathcal{L}_\psi \ : \ P & \to & \mathcal{L}_\psi P \subset \mathbb{R}^n \\ u & = & \mathcal{L}_\psi(x) = \dfrac{\partial \psi}{\partial x}(x). \end{array} \tag{1}$$

This framework gives a new way of obtaining tropical geometry from complex algebraic geometry by degenerating the ambient toric complex structure, while keeping the symplectic manifold structure of X_P unchanged, rather than taking a limit of deformations of the complex field [Mi, EKL].

Another significative difference is that the limit amoebas described above live inside the compact image $\mathcal{L}_\psi P$ and are tropical in the interior of $\mathcal{L}_\psi P$.

Let us describe these results in more detail.

1.1. **Geometric quantization of toric varieties.** Let P be a Delzant polytope with vertices in $\mathbb{Z}^n$ defining, via the Delzant construction [De], a compact symplectic toric manifold $(X_P, \omega, \mathbb{T}^n, \mu_P)$, with moment map μ_P. Let $\mathcal{P}_\mathbb{R} \subset (TX_P)_\mathbb{C}$ be the (singular) real polarization, in the sense of geometric quantization [Wo], corresponding to the orbits of the Hamiltonian $\mathbb{T}^n$ action. The Delzant construction also defines a complex structure J_P on X_P such that the pair (ω, J_P), is Kähler, with Kähler polarization $\mathcal{P}_\mathbb{C}$. In addition, the polytope P defines, canonically, an equivariant J_P-holomorphic line bundle, $L \to X_P$ with curvature $-i\omega$ [Od].

A result, usually attributed to Danilov and Atiyah [Da, GGK], states that the number of integer points in P, which are equal to the images under μ_P of the Bohr-Sommerfeld

(BS) fibers of the real polarization $\mathcal{P}_{\mathbb{R}}$, is equal to the number of holomorphic sections of L, i.e. to the dimensionality of $H^0(X_P, L)$ (see also [Ham1]).

An important general problem in geometric quantization is understanding the relation between quantizations associated to different polarizations and, in particular, between real and holomorphic quantizations. Hitchin [Hi] has shown that, in some general situations, the bundle of quantum Hilbert spaces, over the space of deformations of the complex structure, is equipped with a (projectively) flat connection that provides the identification between holomorphic quantizations corresponding to different complex structures. In [Sn2], Śniatycki has shown that for non-singular real polarizations of arbitrary (quantizable) symplectic manifolds, the set of BS fibers is in bijective correspondence with a generating set for the space of cohomological wave functions which define the quantum Hilbert space in the real polarization. Explicit geometro-analytic relations between real polarization wave functions and holomorphic ones have been found for theta functions on abelian varieties [FMN, BMN], by considering degenerating families of complex structures. For previous related work see [Mu, AdPW, An, Ty, KW]. Similar studies have been performed for cotangent bundles of Lie groups [Hal, FMMN]. Some of the results in this paper, in fact, are related to these results for the case $(T^*\mathbb{S}^1)^n = (\mathbb{C}^*)^n$, where in the present setting $(\mathbb{C}^*)^n$ becomes the open dense orbit in the toric variety X_P.

As opposed to all these cases, however, the real polarization of a compact toric variety always contains singular fibers. As was shown by Hamilton [Ham2], the sheaf cohomology used by Śniatycki only detects the non-singular BS leaves. Also, a possible model for the real quantization that includes the singular fibers has recently been described in [BGU].

If, on one hand, it is natural to expect that by finding a family of (Kähler) complex structures starting at J_P and degenerating to the real polarization, the holomorphic sections will converge to delta distributions supported at the BS fibers, on the other hand, it was unclear how to achieve such behavior from the simple monomial sections characteristic of holomorphic line bundles on toric manifolds (where the series characteristic of theta functions on Abelian varieties are absent). We undertake this study in the present paper for all symplectic toric manifolds and an infinite-dimensional family of toric complex structures giving holomorphic polarizations degenerating to $\mathcal{P}_{\mathbb{R}}$. The study of these degenerations of complex structure in a general symplectic toric manifold is made possible by Abreu's description of toric complex structures [Ab1, Ab2], following Guillemin's characterization of the boundary conditions of the toric data on ∂P [Gui]. For any toric complex structure J on X_P, such that the pair (ω, J) is Kähler, we consider the family of symplectic potentials

$$g_s = g_P + \varphi + s\psi, \tag{2}$$

where g_P is the function defined in (6) [Gui], φ satisfies some regularity conditions (see (7) below and [Ab1, Ab2]), and ψ in the space $C^\infty_+(P)$ of smooth functions with positive definite Hessian on the whole of P. Note that this defines a toric complex structure for every $s \geq 0$ (see (9)).

The quantization of a compact symplectic manifold in the real polarization is given by distributional sections. In this case, conditions of covariant constancy of the wave functions have to be understood as local rather than pointwise (see, for instance, [Ki]) and the relevant piece of data is the sheaf of smooth local sections of a polarization $\mathcal{P}$, $C^\infty(\mathcal{P})$.

Applying this approach just outlined, in Section 3 we obtain a description of the quantum space of the real polarization $\mathcal{Q}_\mathbb{R}$. The main technical difference as compared with the techniques of [Sn2] (applied to the present situation in [Ham2]) consists in the fact that we do not only use the sheaf of sections in the kernel of covariant differentiation, but also the cokernel. It is therefore not surprising that our result differs from that of [Ham2], where the dimension is given by the number of non-degenerate Bohr-Sommerfeld fibers only. In contrast to this, we find

Theorem 1.1. *For the singular real polarization $\mathcal{P}_\mathbb{R}$ defined by the moment map, the space of covariantly constant distributional sections of the prequantum line bundle L_ω is spanned by one section δ^m per Bohr-Sommerfeld fiber $\mu_P^{-1}(m), m \in P \cap \mathbb{Z}^n$, with*

$$\operatorname{supp} \delta^m = \mu_P^{-1}(m).$$

But not only does the result of the quantization in the real polarization change; actually, the weak equations of covariant constancy allow for a continuous passage from quantization in complex to real polarizations. The first step in this direction is to verify that the conditions imposed on distributional sections by the set of equations of covariant constancy converge in a suitable sense: if we denote by $\mathcal{P}^s_\mathbb{C}$ the holomorphic polarization corresponding to the complex structure defined by (2), our second main finding is

Theorem 1.2. *For any $\psi \in C^\infty_+(P)$, we have*

$$C^\infty(\lim_{s\to\infty} \mathcal{P}^s_\mathbb{C}) = C^\infty(\mathcal{P}_\mathbb{R}),$$

where the limit is taken in the positive Lagrangian Grassmannian of the complexified tangent space at each point in X_P.

Identifying holomorphic sections with distributional sections in the usual way (as in [Gun], but making use of the Liouville measure on the base) we may actually keep track of the monomial basis of holomorphic sections as s changes and show that they converge in the space of distributional sections.

Consider the prequantum bundle L_ω equipped with the holomorphic structure defined by the prequantum connection ∇, defined in (13), and by the complex structure J_s corresponding to g_s in (2). Let $\iota : C^\infty(L_\omega) \to (C^\infty(L_\omega^{-1}))'$ be the natural injection of the space of smooth into distributional sections defined in (14). For any lattice point $m \in P \cap \mathbb{Z}^n$, let $\sigma^m_s \in C^\infty(L_\omega)$ be the associated J_s-holomorphic section of L_ω and δ^m the delta distribution from the previous Theorem. Our third main result is the following:

Theorem 1.3. *For any ψ strictly convex in a neighborhood of P and $m \in P \cap \mathbb{Z}^n$, consider the family of L^1-normalized J_s-holomorphic sections*

$$\mathbb{R}^+ \ni s \mapsto \xi^m_s := \frac{\sigma^m_s}{\|\sigma^m_s\|_1} \in C^\infty(L_\omega) \overset{\iota}{\subset} (C^\infty(L_\omega^{-1}))' \quad .$$

Then, as $s \to \infty$, $\iota(\xi^m_s)$ converges to δ^m in $(C^\infty(L_\omega^{-1}))'$.

Remark 1.4. Note that the sections $\sigma^m_s, \sigma^{m'}_s$ are L^2-orthogonal for $m \neq m'$.

Remark 1.5. We note that the set up above can be easily generalized to a larger family of deformations given by symplectic potentials of the form $g_s = g_P + \varphi + \psi_s$, where ψ_s is a family of smooth strictly convex functions on P, such that $\frac{1}{s}\psi_s$ has a strictly convex limit in the C^2-norm in $C^\infty(P)$.

Remark 1.6. For non-compact symplectic toric manifolds X_P, the symplectic potentials in (2) still define compatible complex structures on X_P; however, Abreu's theorem no longer holds. Theorems 1.2 and 1.3 remain valid in the non-compact case, if one assumes uniform strict convexity of ψ for the latter.

As mentioned above, these results provide a setup for relating quantizations in different polarizations. In particular, Theorem 1.3 gives an explicit analytic relation between holomorphic and real wave functions by considering families of complex structures converging to a degenerate point. We will come back to the relation of our results to the more general setup of [Hi, AdPW] in a forthcoming paper.

1.2. **Compact tropical amoebas.** Let now Y_s denote the one-parameter family of hypersurfaces in (X_P, J_s) given by

$$Y_s = \left\{ p \in X_P \ : \ \sum_{m \in P\cap\mathbb{Z}^n} a_m \mathrm{e}^{-sv(m)} \sigma_s^m(p) = 0 \right\}, \tag{3}$$

where $a_m \in \mathbb{C}^*, v(m) \in \mathbb{R}, \forall m \in P \cap \mathbb{Z}^n$. The image of Y_s in P under the moment map μ_P is naturally called the compact amoeba of Y_s. Note that Y_s is a complex submanifold of X_P equipped with the Kähler structure $(J_s, \gamma_s = \omega(.,J_s.))$. This is in contrast with the compact amoeba of [GKZ, FPT, Mi] where the Kähler structure is held fixed.

Using the family of Legendre transforms in (1) associated to the potentials g_s on the open orbit, we relate the intersection $\mu_P(Y_s) \cap \breve{P}$ with the Log_t-amoeba of [FPT, Mi] for finite $t = \mathrm{e}^s$. For $s \to \infty$, the Hausdorff limit of the compact amoeba is then characterized by the tropical amoeba $\mathcal{A}_{\mathrm{trop}}$ defined as the support of non-differentiability, or corner locus, of the piecewise smooth continuous function $\mathbb{R}^n \to \mathbb{R}$,

$$u \ \mapsto \ \max_{m \in P\cap\mathbb{Z}^n} \left\{ {}^t m \cdot u - v(m) \right\}.$$

In Section 6, we show that, as $s \to \infty$, the amoebas $\mu_P(Y_s)$ converge in the Hausdorff metric to a limit amoeba $\mathcal{A}_{\mathrm{lim}}$. The relation between $\mathcal{A}_{\mathrm{trop}}$ and $\mathcal{A}_{\mathrm{lim}}$ is given by a projection $\pi : \mathbb{R}^n \to \mathcal{L}_\psi P$ (defined in Lemma 6.7, see Figure 1) determined by ψ and the combinatorics of the fan of P. The fourth main result is, then:

Theorem 1.7. *The limit amoeba is given by*

$$\mathcal{L}_\psi \mathcal{A}_{\mathrm{lim}} = \pi \mathcal{A}_{\mathrm{trop}}.$$

Remark 1.8. For an open set of valuations $v(m)$ in (3), the convex projection $\pi\mathcal{A}_{\mathrm{trop}}$ coincides with the intersection of $\mathcal{A}_{\mathrm{trop}}$ with the image of the moment polytope $\mathcal{L}_\psi P$. In particular, if $\psi(x) = \frac{x^2}{2}$, then $\mathcal{L}_\psi = \mathrm{Id}_P$ and $\mathcal{A}_{\mathrm{lim}}$ is a (compact part) of a tropical amoeba, see Figure 3.

Remark 1.9. Note that for quadratic $\psi(x) = \frac{{}^t x G x}{2} + {}^t b x$, where ${}^t G = G > 0$, the limit amoeba $\mathcal{A}_{\mathrm{lim}} \subset P$ itself is piecewise linear.

This construction produces naturally a singular affine manifold $\mathcal{L}_\psi \mathcal{A}_{\mathrm{lim}}$, with a metric structure (induced from the inverse of the Hessian of ψ). In the last Section we comment on the possible relation of this result to the study of large complex structure limits in homological mirror symmetry.

2. Preliminaries and notation

Let us briefly review a few facts concerning compatible complex structures on toric symplectic manifolds and also fix some notation. For reviews on toric varieties, see [Co, Da, dS]. Consider a Delzant lattice polytope $P \subset \mathbb{R}^n$ given by

$$P = \{x \in \mathbb{R}^n \ : \ \ell_r(x) \geq 0, r = 1, \dots, d\}, \tag{4}$$

where

$$\begin{aligned} \ell_r \ : \ \mathbb{R}^n & \rightarrow \mathbb{R} \\ \ell_r(x) & = {}^t\nu_r x - \lambda_r, \end{aligned} \tag{5}$$

$\lambda_r \in \mathbb{Z}$ and ν_r are primitive vectors of the lattice $\mathbb{Z}^n \subset \mathbb{R}^n$, inward-pointing and normal to the r-th facet, i.e. codimension-1 face of P. Let $\breve{P}$ be the interior of P.

Let X_P be the associated smooth toric variety, with moment map $\mu_P : X_P \rightarrow P$. On the open dense orbit $\breve{X}_P = \mu_P^{-1}(\breve{P}) \cong \breve{P} \times \mathbb{T}^n$, one considers symplectic, or action-angle, coordinates $(x, \theta) \in \breve{P} \times \mathbb{T}^n$ for which the symplectic form is the standard one, $\omega = \sum_{i=1}^n dx_i \wedge d\theta_i$ and $\mu_P(x, \theta) = x$.

Recall ([Ab1, Ab2]) that any compatible complex structure, J, on X_P can be written via a symplectic potential $g = g_P + \varphi$, where $g_P \in C^\infty(\breve{P})$ is given by [Gui]

$$g_P(x) = \frac{1}{2} \sum_{r=1}^{d} \ell_r(x) \log \ell_r(x), \tag{6}$$

and $\varphi \in C^\infty_{g_P}(P)$ belongs to the convex set, $C^\infty_{g_P}(P) \subset C^\infty(P)$, of functions φ such that $\mathrm{Hess}_x(g_P + \varphi)$ is positive definite on $\breve{P}$ and satisfies the regularity conditions,

$$\det(\mathrm{Hess}_x(g_P + \varphi)) = \left[\alpha(x) \Pi_{r=1}^d \ell_r(x)\right]^{-1}, \tag{7}$$

where α is smooth and strictly positive on P, as described in [Ab1, Ab2]. Let $C^\infty_+(P) \subset C^\infty_{g_P}(P) \subset C^\infty(P)$ be the convex set of functions with positive definite Hessian on the whole of P. Note that the function

$$g_s = g_P + \varphi + s\psi, \tag{8}$$

where $\varphi \in C^\infty_{g_P}(P), \psi \in C^\infty_+(P)$ defines a toric complex structure for every $s \geq 0$ (see (9)).

The complex structure J and the metric $\gamma = \omega(\cdot, J\cdot)$ are given by

$$J = \begin{pmatrix} 0 & -G^{-1} \\ G & 0 \end{pmatrix} \ ; \ \gamma = \begin{pmatrix} G & 0 \\ 0 & G^{-1} \end{pmatrix}, \tag{9}$$

where $G = \mathrm{Hess}_x g$ is the Hessian of g. Note that g is strictly convex. For recent applications of this result see e.g. [Do, MSY, SeD].

The complex coordinates are related with the symplectic ones by a bijective Legendre transform

$$\breve{P} \ni x \mapsto y = \frac{\partial g}{\partial x} \in \mathbb{R}^n,$$

that is, g fixes an equivariant biholomorphism $\breve{P}\times\mathbb{T}^n\cong(\mathbb{C}^*)^n$,

$$\breve{P}\times\mathbb{T}^n\ni(x,\theta)\mapsto w=\mathrm{e}^{y+\mathrm{i}\theta}\in(\mathbb{C}^*)^n.$$

The inverse transformation is given by $x=\frac{\partial h}{\partial y}$, where

$$h(y)={}^t x(y)y-g(x(y)). \tag{10}$$

Let us describe coordinate charts covering the rest of X_P, that is the loci of compactification. Using the Legendre transform associated to the symplectic potential g, one can describe, in particular, holomorphic charts around the fixed points of the torus action. Consider, for any vertex v of P, any ordering of the n facets that contain v; upon reordering the indices, one may suppose that

$$\ell_1(v)=\cdots=\ell_n(v)=0.$$

Consider the affine change of variables on P

$$l_i=\ell_i(x)={}^t\nu_i x-\lambda_i,\quad\forall i=1,\dots,n,\ \text{i.e.}\quad l=Ax-\lambda,$$

where $A=(A_{ij}=(\nu_i)_j)\in Gl(n,\mathbb{Z})$ and $\lambda\in\mathbb{Z}^n$. This induces a change of variables on the open orbit $\breve{X}_P$,

$$\breve{P}\times\mathbb{T}^n\ni(x,\theta)\mapsto(l=Ax-\lambda,\vartheta={}^tA^{-1}\theta)\in(AP-\lambda)\times\mathbb{T}^n\subset(\mathbb{R}_0^+)^n\times\mathbb{T}^n$$

such that $\omega=\sum\mathrm{d}x_i\wedge\mathrm{d}\theta_i=\sum\mathrm{d}l_i\wedge\mathrm{d}\vartheta_i$.

Consider now the union

$$\breve{P}_v:=\{v\}\cup\bigcup_{\substack{F\text{ face}\\ v\in\overline{F}}}\breve{F}$$

of the interior of all faces adjacent to v and set $V_v:=\mu_P^{-1}(\breve{P}_v)\subset X_P$. This is an open neighborhood of the fixed point $\mu_P^{-1}(v)$, and it carries a smooth chart $V_v\to\mathbb{C}^n$ that glues to the chart on the open orbit as

$$(A\breve{P}-\lambda)\times\mathbb{T}^n\ni(l,\vartheta)\mapsto w_v=(w_{l_j}=\mathrm{e}^{y_{l_j}+\mathrm{i}\vartheta_j})_{j=i}^n\in\mathbb{C}^n, \tag{11}$$

where $y_{l_j}=\frac{\partial g}{\partial l_j}$. We will call this "the chart at v" for short, dropping any reference to the choice of ordering of the facets at v to disburden the notation; usually we will then write simply w_j for the components of w_v. It is easy to see that a change of coordinates between two such charts, at two vertices v and $\widetilde{v}$, is holomorphic. The complex manifold, W_P, obtained by taking the vertex complex charts and the transition functions between them, does not depend on the symplectic potential. It will be convenient for us to distinguish between X_P and W_P, noting that the g-dependent map $\chi_g:X_P\to W_P$ described locally by (11), introduces the g-dependent complex structure (9) on X_P, making χ_g a biholomorphism. In the same way, the holomorphic line bundle L_P on W_P determined canonically by the polytope P [Od], induces, via the pull-back by χ_g, an holomorphic structure on the (smooth) prequantum line bundle L_ω on X_P, as will be described below:

$$\begin{tikzcd}
L_\omega \arrow[r] \arrow[d] & L_P \arrow[d] \\
X_P \arrow[r, "\chi_g"] & W_P
\end{tikzcd}.$$

Using the charts V_v at the fixed points, one can describe L_P by

$$L_P = \left(\coprod_v V_v \times \mathbb{C} \right) / \sim,$$

where the equivalence relation $\sim$ is given by the transition functions for the local trivializing holomorphic sections $\mathbb{1}_v(w_v) = (w_v, 1)$ for $p \in V_v$,

$$\mathbb{1}_v = w_{\widetilde{v}}^{(\widetilde{A}A^{-1}\lambda - \widetilde{\lambda})} \mathbb{1}_{\tilde{v}},$$

on intersections of the domains. Here, we use the data of the affine changes of coordinates

$$l = \ell(x) = Ax - \lambda, \qquad \widetilde{l} = \widetilde{\ell}(x) = \widetilde{A}x - \widetilde{\lambda},$$

associated to the vertices v and $\widetilde{v}$ (and the order of the facets there).

Note that one also has a trivializing section on the open orbit, such that for a vertex $v \in P$, $\mathbb{1}_v = w^v \breve{\mathbb{1}}$ on $\breve{W}_P$. However, note that the sections $\mathbb{1}_v$ extend to global holomorphic sections on W_P while the extension of $\breve{\mathbb{1}}$ will, in general, be meromorphic and will be holomorphic iff $0 \in P \cap \mathbb{Z}^n$. Sections in the standard basis $\{\sigma^m\}_{m \in P \cap \mathbb{Z}^n} \subset H^0(W_P, L_P)$ read $\sigma^m = \sigma^m_{\breve{P}} \breve{\mathbb{1}} = \sigma^m_v \mathbb{1}_v$, with

$$\sigma^m_{\breve{P}}(w) = w^m, \qquad \sigma^m_v(w_v) = w_v^{\ell(m)},$$

in the respective domains.

The prequantum line bundle L_ω on the symplectic manifold X_P is, analogously, defined by unitary local trivializing sections $\mathbb{1}_v^{U(1)}$ and transition functions

$$\mathbb{1}_v^{U(1)} = \mathrm{e}^{\mathrm{i}\,{}^t(\widetilde{A}A^{-1}\lambda - \widetilde{\lambda})\tilde{\vartheta}} \mathbb{1}_{\tilde{v}}^{U(1)}. \tag{12}$$

L_ω is equipped with the compatible prequantum connection ∇, of curvature $-\mathrm{i}\omega$, defined by

$$\nabla \mathbb{1}_v^{U(1)} = -\mathrm{i}\,{}^t(x - v)\mathrm{d}\theta\, \mathbb{1}_v^{U(1)} = -\mathrm{i}\,{}^t l\, \mathrm{d}\vartheta\, \mathbb{1}_v^{U(1)}, \tag{13}$$

where we use $\ell(v) = 0$.

The bundle isomorphism relating L_ω and L_P is determined by

$$\mathbb{1}_v^{U(1)} = \mathrm{e}^{h_v \circ \mu_P} \chi_g^* \mathbb{1}_v, \qquad \breve{\mathbb{1}}^{U(1)} = \mathrm{e}^{h \circ \mu_P} \chi_g^* \breve{\mathbb{1}},$$

where, for $m \in \mathbb{Z}^n$, $h_m(x) = (x - m)\frac{\partial g}{\partial x} - g(x)$ and h is the function in (10) defining the inverse Legendre transform.

In these unitary local trivializations, the sections $\chi_g^* \sigma^m$ read

$$\chi_g^* \sigma^m = \mathrm{e}^{-h_m \circ \mu_P} \mathrm{e}^{\mathrm{i}\,{}^t m \theta} \breve{\mathbb{1}}^{U(1)} = \mathrm{e}^{-h_m \circ \mu_P} \mathrm{e}^{\mathrm{i}\,{}^t \ell(m) \vartheta} \mathbb{1}_v^{U(1)},$$

where, after an affine change of coordinates $x \mapsto \ell(x)$ on the moment polytope, as above, we get $h_m(l) = {}^t(l - \ell(m))\frac{\partial g}{\partial l} - g(l)$. Then, for all $\sigma \in H^0(W_P, L_P)$, $\chi_g^* \sigma \in C^\infty(L_\omega)$ is holomorphic, that is

$$\nabla_{\overline{\xi}} \chi_g^* \sigma = 0,$$

for any holomorphic vector field ξ. That is, such sections are polarized with respect to the distribution of holomorphic vector fields on X_P (see, for instance, [Wo]).

It is critical to our approach to pass from the sheaf of smooth sections of L_ω to the sheaf of distributional sections: using the fixed Liouville measure ω^n on X_P, we can identify the topological dual of the locally convex vector space of smooth sections of

L_ω^{-1} on X_P, $C^\infty(L_\omega^{-1})$, with distributional sections of the bundle L_ω (see, for instance, [Gun]):

$$\begin{aligned} \iota \quad &: \quad C^\infty(L_\omega) \hookrightarrow (C^\infty(L_\omega^{-1}))' \\ \iota\sigma(\tau) \quad &= \quad \int_{X_P} \sigma\tau\omega^n, \qquad \forall \tau \in C^\infty(L_\omega^{-1}). \end{aligned} \tag{14}$$

In the presence of the unitary structure on L_ω given in (12), $\overline{L_\omega}^{-1}$ and L_ω are naturally identified, and therefore

$$\iota\sigma(\tau) = \langle \sigma, \overline{\tau} \rangle_{L^2}.$$

Let ∇_ξ^* be the adjoint of the (unbounded) operator ∇_ξ on the Hilbert space $L^2(L_\omega)$. Then, one has

$$\iota(\nabla_\xi\sigma)(\tau) = \langle \nabla_\xi\sigma, \overline{\tau} \rangle_{L^2} = \langle \sigma, \overline{\overline{\nabla_\xi^*}\tau} \rangle_{L^2},$$

where $\overline{\nabla}^*$ is the connection induced by ∇^* on $\overline{L_\omega}$, that is

$$\iota(\nabla_\xi\sigma)(\tau) = \int_{\check{X}_P} (\mathrm{d}_\xi - \mathrm{i}^t x \mathrm{d}\theta \cdot \xi)(\sigma)\tau\omega^n = -\int_{\check{X}_P} \sigma(\mathrm{d}_\xi + \mathrm{i}^t x \mathrm{d}\theta \cdot \xi)(\tau)\omega^n. \tag{15}$$

Covariant differentiation of distributional sections is then defined by

$$(\nabla''_\xi \eta)(\tau) := \eta(\overline{\nabla_\xi^*}\tau), \quad \forall \tau \in C^\infty(L^{-1}),$$

so that, naturally,

$$\iota(\nabla_\xi\sigma) = \nabla''_\xi(\iota\sigma) \quad \forall \sigma \in C^\infty(L_\omega).$$

Remark 2.1. Note that in the same way, covariant differentiation ∇'' is defined by integration by parts on arbitrary open sets $W \subset X_P$. Naturally, the sheaf of distributional sections of L_ω over an open set W has sections

$$C^{-\infty}(L_\omega|_W) = \left(C_c^\infty(L_\omega^{-1}|_W)\right)'$$

that must be tested against smooth sections with compact support.

3. QUANTIZATION IN A REAL POLARIZATION

In this Section we describe the quantization obtained using the definition outlined above for the singular real polarization $\mathcal{P}_\mathbb{R}$ defined by the moment map

$$\mathcal{P}_\mathbb{R} = \ker \mathrm{d}\mu_P.$$

This means that we consider the space of weakly covariant constant distributional sections, $\mathcal{Q}_\mathbb{R} \subset (C^\infty(L_\omega^{-1}))'$,

$$\mathcal{Q}_\mathbb{R} := \left\{\sigma \in (C^\infty(L_\omega^{-1}))' | \forall\, W \subset X_P \text{ open}, \forall\, \xi \in C^\infty(\mathcal{P}_\mathbb{R}|_W),\ \nabla''_\xi(\sigma|_W) = 0\right\}.$$

In the present Section, "covariantly constant" is always understood to mean "covariantly constant with respect to the polarization $\mathcal{P}_\mathbb{R}$". We first give a result describing the local covariantly constant sections. Note that it only depends on the local structure of the polarization, and thus applies also in cases where globally one is not dealing with toric varieties.

Proposition 3.1. (i) *Any covariantly constant section on a $\mathbb{T}^n$-invariant open set $W \subset X_P$ is supported on the Bohr-Sommerfeld fibers in W.*

(ii) *If W contains the Bohr-Sommerfeld fiber $\mu_P^{-1}(m)$ (with $m \in P \cap \mathbb{Z}^n$), the distributional section*

$$\delta^m(\tau) = \int\limits_{\mu_P^{-1}(m)} \mathrm{e}^{\mathrm{i}\,{}^t\ell(m)\vartheta}\tau_v, \quad \forall \tau \in C_c^\infty(L_\omega^{-1}|_W),$$

(where integration is understood with respect to Haar measure on the torus $\mu_P^{-1}(m)$, $v \in P$ is a vertex such that $m \in V_v$ and $\tau = \tau_v \mathbb{1}_v^{U(1)}$ on V_v) is covariantly constant.

(iii) *The sections δ^m, $m \in \mu_P(W) \cap \mathbb{Z}^n$ span the space of covariantly constant sections on W.*

Proof. Note that all the statements are local in nature. Using the action of $Sl(n, \mathbb{Z})$ on polytopes, we can reduce without loss of generality an arbitrary Bohr-Sommerfeld fiber $\mu_P^{-1}(m)$ with $m \in P \cap \mathbb{Z}^n$ being contained in a codimension k (and no codimension $k+1$) face of P to the form

$$m = (\underbrace{0, \ldots, 0}_{k}, \widetilde{m}) \qquad \text{with } \widetilde{m}_j > 0 \quad \forall j = k+1, \ldots, n.$$

Furthermore, we can cover all of X_P by charts $\widetilde{W}$ of special neighborhoods of $\mu_P^{-1}(m)$ of the form

$$\widetilde{W} := B_\varepsilon(0) \times \left(\widetilde{m} +]-\varepsilon, \varepsilon[^{n-k}\right) \times \mathbb{T}^{n-k}$$

with $0 < \varepsilon < 1$ and $B_\varepsilon(0) \subset \mathbb{R}^{2k}$ being the ball of radius ε in $\mathbb{R}^{2k}$. This chart is glued onto the open orbit via the map,

$$\breve{P} \times \mathbb{T}^n \ni (x_1, \ldots, x_n, \vartheta_1, \ldots, \vartheta_n) \mapsto$$
$$\big(u_1 = \sqrt{x_1}\cos\vartheta_1, v_1 = \sqrt{x_1}\sin\vartheta_1, \ldots, u_k = \sqrt{x_k}\cos\vartheta_k, v_k = \sqrt{x_k}\sin\vartheta_k,$$
$$x_{k+1}, \ldots, x_n, \vartheta_{k+1}, \ldots, \vartheta_n) \in \widetilde{W}$$

and therefore induces the standard symplectic form in the coordinates u, v, x, ϑ

$$\omega = \sum_{j=1}^{k} \mathrm{d}u_j \wedge \mathrm{d}v_j + \sum_{j=k+1}^{n} \mathrm{d}x_j \wedge \mathrm{d}\vartheta_j.$$

We then trivialize the line bundle with connection L_ω over $\widetilde{W}$ by setting

$$\nabla = \mathrm{d} - \mathrm{i}\left(\frac{1}{2}({}^t u\mathrm{d}v - {}^t v\mathrm{d}u) + {}^t x\mathrm{d}\vartheta\right).$$

Therefore, we are reduced to studying the equations of covariant constancy on the space of (usual) distributions $C^{-\infty}(\widetilde{W})$ on $\widetilde{W}$, using arbitrary test functions in $C_c^\infty(\widetilde{W})$ and vector fields $\xi \in C^\infty(\mathcal{P}_\mathbb{R}|_{\widetilde{W}})$. These can be written as

$$\xi_{(u,v,x,\vartheta)} = \sum_{j=1}^{k} \alpha_j \left(u_j \frac{\partial}{\partial v_j} - v_j \frac{\partial}{\partial u_j}\right) + \sum_{j=k+1}^{n} \beta_j \frac{\partial}{\partial \vartheta_j}, \tag{16}$$

where α_j, β_j are smooth functions on $\widetilde{W}$.

(i) First, note that if a distribution σ is covariantly constant, it is unchanged by parallel transport around a loop in $\mathbb{T}^n$, while on the other hand this parallel transport results in multiplication of all test sections by the holonomies of the respective leaves. Explicitly, for a loop specified by a vector $a \in \mathbb{Z}^n$, any test function $\tau \in C_c^\infty(\widetilde{W})$ is multiplied by the smooth function $\mathrm{e}^{2\pi\mathrm{i}^t a\, \mu_P(u,v,x,\vartheta)}$. Therefore,

$$\sigma = \mathrm{e}^{2\pi\mathrm{i}^t a\, \mu_P}\sigma, \quad \forall a \in \mathbb{Z}^n,$$

and σ must be supported in the set where all the exponentials equal 1, i.e. on the Bohr-Sommerfeld fibers, where μ_P takes integer values.

(ii) In the chart on $\widetilde{W}$, any test function τ can be written as a Fourier series

$$\tau(u,v,x,\vartheta) = \sum_{b\in\mathbb{Z}^{n-k}} \widehat{\tau}(u,v,x,b)\mathrm{e}^{\mathrm{i}^t b\,\vartheta},$$

with coefficients that are smooth and compactly supported in the other variables,

$$\forall b \in \mathbb{Z}^{n-k}: \quad \widehat{\tau}(.,b) \in C_c^\infty\left(B_\varepsilon(0) \times \left(\widetilde{m} +]-\varepsilon,\varepsilon[^{n-k}\right)\right).$$

The distributional section δ^m is calculated as

$$\begin{aligned} \delta^m(\tau) &= \int\limits_{\mu_P^{-1}(m)} \mathrm{e}^{\mathrm{i}^t\widetilde{m}\vartheta}\tau = \\ &= \int\limits_{\mathbb{T}^{n-k}} \mathrm{e}^{\mathrm{i}^t\widetilde{m}\vartheta}\tau(u=0,v=0,x=\widetilde{m},\vartheta)\mathrm{d}\vartheta = \\ &= \widehat{\tau}(u=0,v=0,x=\widetilde{m},b=-\widetilde{m}). \end{aligned}$$

We can write this in abreviated form as

$$\delta^m(\tau) = \widehat{\tau}(m,-m),$$

where

$$\widehat{\tau}(x,c) = \int\limits_{\mu_P^{-1}(x)} \mathrm{e}^{\mathrm{i}\ell(-c)\vartheta}\tau_v, \quad c \in \mathbb{Z}^n, \tag{17}$$

is the fiberwise Fourier transform of τ on the open orbit, evaluated for $c = -m$ in the limit $x \to m$.

Differentiating an arbitrary test function τ along an arbitrary vector field ξ as in equation (16) gives

$$\begin{aligned} \overline{\nabla}^*_\xi\tau &= -\mathrm{d}\tau\xi - \mathrm{i}\left(\frac{1}{2}({}^tu\mathrm{d}v - {}^tv\mathrm{d}u) + {}^tx\mathrm{d}\vartheta\right)\xi\tau = \\ &= -\sum_{b\in\mathbb{Z}^{n-k}}\left(\sum_{j=1}^k \alpha_j\left(u_j\frac{\partial\widehat{\tau}}{\partial v_j} - v_j\frac{\partial\widehat{\tau}}{\partial u_j} + \frac{\mathrm{i}}{2}(u_j^2+v_j^2)\widehat{\tau}\right) + \right. \\ &\qquad \left. +\mathrm{i}\sum_{j=k+1}^n \beta_j(x_j+b_j)\widehat{\tau}\right)\mathrm{e}^{\mathrm{i}^t b\,\vartheta}, \end{aligned} \tag{18}$$

whence for arbitrary τ and ξ

$$\begin{aligned} \delta^m(\overline{\nabla}^*_\xi\tau) &= \left(\sum_{j=1}^k \alpha_j\left(u_j\frac{\partial\widehat{\tau}}{\partial v_j} - v_j\frac{\partial\widehat{\tau}}{\partial u_j} + \frac{\mathrm{i}}{2}(u_j^2+v_j^2)\widehat{\tau}\right) + \right. \\ &\qquad \left. +\mathrm{i}\sum_{j=k+1}^n \beta_j(x_j+b_j)\widehat{\tau}\right)_{(u=0,v=0,x=\widetilde{m},b=-\widetilde{m})} = 0, \end{aligned}$$

so that δ^m is covariantly constant.

(iii) Consider an arbitrary distribution $\sigma \in C^{-\infty}(\widetilde{W})$. Using Fourier expansion of test functions along the non-degenerating directions of the polarization on $\widetilde{W}$, as in item (ii), to σ we associate a family of distributions $\widehat{\sigma}_b$ on $B_\varepsilon(0) \times \left(\widetilde{m} +]-\varepsilon, \varepsilon[^{n-k}\right)$ by setting

$$\widehat{\sigma}_b(\psi) := \sigma\left(\psi(u,v,x)\mathrm{e}^{\mathrm{i}\,{}^t b\vartheta}\right), \quad \forall \psi \in C_c^\infty\left(B_\varepsilon(0) \times \left(\widetilde{m} +]-\varepsilon, \varepsilon[^{n-k}\right)\right),$$

so that

$$\sigma(\tau) = \sum_{b \in \mathbb{Z}^{n-k}} \widehat{\sigma}_b(\widehat{\tau}(.,b)).$$

It is clear that the map $\sigma \mapsto (\widehat{\sigma}_b)_{b \in \mathbb{Z}^{n-k}}$ is injective, so we need to show that the condition of covariant constancy

$$\sigma(\overline{\nabla}_\xi^* \tau) = 0, \quad \forall \tau \in C_c^\infty(\widetilde{W}), \xi \in C^\infty(\mathcal{P}_\mathbb{R}|_{\widetilde{W}})$$

implies that

$$\widehat{\sigma}_b = \begin{cases} 0 & \text{if } b \neq -\widetilde{m}, \\ \lambda\delta(u,v,x-\widetilde{m}) & \text{if } b = -\widetilde{m}, \text{ where } \lambda \in \mathbb{C}. \end{cases}$$

From (i) we know that $\widehat{\sigma}_b$ has support at the point $(u,v,x) = (0,0,\widetilde{m})$ for each b, therefore it must be of the form (see, for instance, [Tr] Chapter 24; here j,k,l are multi-indices)

$$\widehat{\sigma}_b = \sum_{\text{finite}} \gamma_{jkl}^b \frac{\partial^{|j|+|k|+|l|}}{\partial u^j \partial v^k \partial x^l} \delta(u,v,x-\widetilde{m}).$$

Using for example a vector field ξ with $\alpha_j = \beta_j \equiv 1$ and explicit test functions τ that are polynomial near the point $(u,v,x) = (0,0,\widetilde{m})$, it is easy to give examples that show that if $\gamma_{jkl}^b \neq 0$ and either $b \neq -\widetilde{m}$ or $|j|+|k|+|l| > 0$, then there exist ξ and τ such that

$$\sigma(\overline{\nabla}_\xi^* \tau) \neq 0,$$

thus producing a contradiction. □

Immediately from this proposition follows the

***Proof of Theorem* 1.1.** Since the open neighborhoods considered in the Proposition are $\mathbb{T}^n$-invariant and the covariantly constant sections are supported on closed subsets, each of them can evidently be extended by 0 to give a global covariantly constant section. □

4. LARGE TORIC COMPLEX STRUCTURE LIMITS OF KÄHLER POLARIZATIONS

As mentioned above, we will turn our attention to a family of compatible complex structures determined by the symplectic potentials $g_s = g_P + \varphi + s\psi$, being interested in the limit of holomorphic polarizations, in the sense of geometric quantization, and subsequently in the convergence of monomial sections to certain delta distributions.

In the vertex coordinate charts V_v described above, the holomorphic polarizations are given by

$$\mathcal{P}_\mathbb{C}^s = \mathrm{span}_\mathbb{C}\left\{\frac{\partial}{\partial w_j^s} : j = 1,\dots,n\right\}.$$

Let $\mathcal{P}_{\mathbb{R}}$ stand for the vertical polarization, that is

$$\mathcal{P}_{\mathbb{R}} := \ker \mathrm{d}\mu_P,$$

which is real and singular above the boundary ∂P. Let now $\mathcal{P}^\infty = \lim_{s\to\infty} \mathcal{P}^s_{\mathbb{C}}$ where the limit is taken in the positive Lagrangian Grassmannian of the complexified tangent space at each point in X_P.

Lemma 4.1. *On the open orbit* $\check{X}_P$, $\mathcal{P}^\infty = \mathcal{P}_{\mathbb{R}}$.

Proof. By direct calculation,

$$\frac{\partial}{\partial y^s_{l_j}} = \sum_k (G_s^{-1})_{jk} \frac{\partial}{\partial l_k}$$

where

$$(G_s)_{jk} = \operatorname{Hess} g_s$$

is the Hessian of g_s. Since $G_s > s\operatorname{Hess}\psi > 0$, $(G_s^{-1})_{jk} \to 0$ and

$$\operatorname{span}_{\mathbb{C}} \{\frac{\partial}{\partial w^s_j}\} = \operatorname{span}_{\mathbb{C}} \{\frac{\partial}{\partial y^s_{l_j}} - \mathrm{i}\frac{\partial}{\partial \vartheta_j}\} \to \operatorname{span}_{\mathbb{C}} \{\frac{\partial}{\partial \vartheta_j}\}.$$

□

At the points of X_P that do not lie in the open orbit, the holomorphic polarization "in the degenerate angular directions" is independent of g_s, in the following sense:

Lemma 4.2. *Consider two charts around a fixed point* $v \in P$, $w_v, \widetilde{w}_v : V_v \to \mathbb{C}^n$, *specified by symplectic potentials* $g \neq \widetilde{g}$.

Whenever $w_j = 0$, *also* $\widetilde{w}_j = 0$, *and at these points*

$$\mathbb{C} \cdot \frac{\partial}{\partial w_j} = \mathbb{C} \cdot \frac{\partial}{\partial \widetilde{w}_j}, \quad j = 1, \ldots, n.$$

Proof. According to the description of the charts,

$$w_j = \widetilde{w}_j f,$$

where f is a real-valued function, smooth in P, that factorizes through μ_P, that is through $|\widetilde{w}_1|, \ldots, |\widetilde{w}_n|$. Therefore,

$$\mathrm{d}w_j = \sum_k \left[(\delta_{j,k} f + \widetilde{w}_j \frac{\partial f}{\partial \widetilde{w}_k})\mathrm{d}\widetilde{w}_k + \widetilde{w}_j \frac{\partial f}{\partial \overline{\widetilde{w}}_k} \mathrm{d}\overline{\widetilde{w}}_k\right]$$

and at point with $\widetilde{w}_j = 0$ one finds, in fact,

$$\mathbb{C} \cdot \frac{\partial}{\partial w_j} = \mathbb{C} \cdot \frac{\partial}{\partial \widetilde{w}_j}.$$

□

The two lemmata together give

Theorem 4.3. *In any of the charts* w_v, *the limit polarization* $\mathcal{P}^\infty$ *is*

$$\mathcal{P}^\infty = \mathcal{P}_{\mathbb{R}} \oplus \operatorname{span}_{\mathbb{C}} \{\frac{\partial}{\partial w_j} : w_j = 0\}$$

Proof. It suffices to show the convergence

$$\operatorname{span}_{\mathbb{C}} \{\frac{\partial}{\partial w^s_k}\} \to \operatorname{span}_{\mathbb{C}} \{\frac{\partial}{\partial \theta_k}\}$$

whenever $w_k \neq 0$, which really is a small modification of Lemma 4.1. For any face F in the coordinate neighborhood, we write abusively $j \in F$ if $w_j = 0$ along F. For any such affine subspace we have then

$$\begin{aligned} \mathcal{P}^s_{\mathbb{C}} &= \operatorname{span}_{\mathbb{C}} \{\frac{\partial}{\partial w^s_j} : j \in F\} \oplus \operatorname{span}_{\mathbb{C}} \{\frac{\partial}{\partial y^s_{l_k}} - \mathrm{i}\frac{\partial}{\partial \vartheta_k} : k \notin F\} = \\ &= \operatorname{span}_{\mathbb{C}} \{\frac{\partial}{\partial w^s_j} : j \in F\} \oplus \operatorname{span}_{\mathbb{C}} \{\sum_{k' \notin F} ((G_s)_F)^{-1}_{kk'} \frac{\partial}{\partial l_{k'}} - \mathrm{i}\frac{\partial}{\partial \vartheta_k} : k \notin F\} \end{aligned}$$

where $(G_s)_F$ is the minor of G_s specified by the variables that are unrestricted along F, which is well-defined and equals the Hessian of the restriction of g_s there. Since this tends to infinity, its inverse goes to zero and the statement follows. $\square$

This results in

***Proof of Theorem* 1.2.** Given Theorem 4.3, we are reduced to proving that

$$C^\infty\left(\mathcal{P}_{\mathbb{R}} \oplus \operatorname{span}_{\mathbb{C}} \{\frac{\partial}{\partial w_j} : w_j = 0\}\right) = C^\infty(\mathcal{P}_{\mathbb{R}}).$$

This is clear since any smooth complexified vector field $\xi \in C^\infty(T^{\mathbb{C}}X)$ that restricts to a section of $\mathcal{P}_{\mathbb{R}}$ on an open dense subset must satisfy $\overline{\xi} = \xi$ throughout. Such a vector field cannot have components along the directions of $\operatorname{span}_{\mathbb{C}} \{\frac{\partial}{\partial w_j} : w_j = 0\}$. $\square$

Remark 4.4. Note that the Cauchy-Riemann conditions hold for distributions (see e.g. [Gun] for the case $n = 1$, or Lemma 2 in [KY]), that is, for any complex polarization $\mathcal{P}_{\mathbb{C}}$, considering the intersection of the kernels of $\nabla''_{\frac{\partial}{\partial \bar{z}_j}}$ gives exactly the space

$$\bigcap_{\frac{\partial}{\partial \bar{z}_j} \in \overline{\mathcal{P}_{\mathbb{C}}}} \ker \nabla''_{\frac{\partial}{\partial \bar{z}_j}} = \iota H^0(X_P(\mathcal{P}_{\mathbb{C}}), L_\omega(\mathcal{P}_{\mathbb{C}})) \subset (C^\infty(L_\omega^{-1}))'$$

of holomorphic sections (viewed as distributions). Thus one can view the 1-parameter family of quantizations associated to g_s and the real quantization on equal footing, embedded in the space of distributional sections, $\mathcal{Q}_s \subset (C^\infty(L_\omega^{-1}))'$,

$$\mathcal{Q}_s := \left\{\sigma \in (C^\infty(L_\omega^{-1}))' | \forall\, W \subset X_P \text{ open}, \forall\, \xi \in C^\infty(\overline{\mathcal{P}^s}|_W), \nabla''_\xi(\sigma|_W) = 0\right\},$$

for $s \in [0, \infty]$, where $\mathcal{P}^\infty := \mathcal{P}_{\mathbb{R}}$, and $\mathcal{Q}_\infty = \mathcal{Q}_{\mathbb{R}}$ in the previous notation. In the next Section, we will see that the convergence of polarizations proved here translates eventually into a continuous variation of the subspace $\mathcal{Q}_s$ in the space of distributional sections as $s \to \infty$.

Remark 4.5. Notice that Theorem 1.2 in this and Proposition 3.1 in the previous Section are also valid for non-compact P, with some obvious changes such as taking test sections with compact support.

5. LARGE COMPLEX STRUCTURE LIMITS OF HOLOMORPHIC SECTIONS AND BS FIBERS

Here, we use convexity to show that, as $s \to \infty$, the holomorphic sections converge, when normalized properly, to the distributional sections δ^m, described in Proposition 3.1, and that are supported along the Bohr-Sommerfeld fibers of μ_P and covariantly constant along the real polarization.

First, we show an elementary lemma on certain Dirac sequences associated with the convex function ψ that will permit us to prove Theorem 1.3.

Lemma 5.1. *For any ψ strictly convex in a neighborhood of the moment polytope P and any $m \in P \cap \mathbb{Z}^n$, the function*

$$P \ni x \mapsto f_m(x) := {}^t(x-m)\frac{\partial \psi}{\partial x} - \psi(x) \in \mathbb{R}$$

has a unique minimum at $x = m$ and

$$\lim_{s\to\infty} \frac{\mathrm{e}^{-sf_m}}{\|\mathrm{e}^{-sf_m}\|_1} \to \delta(x-m),$$

in the sense of distributions.

Proof. For x in a convex neighborhood of P, using $f_m(m) = -\psi(m)$ and $\nabla f_m(x) = {}^t(x-m)\mathrm{Hess}_x\psi$, we have

$$\begin{aligned} f_m(x) &= f_m(m) + \int_0^1 \frac{d}{dt} f_m(m+t(x-m))\mathrm{d}t = \\ &= -\psi(m) + \int_0^1 {}^t(x-m)\mathrm{Hess}_{m+t(x-m)}\psi(x-m)\,t\,\mathrm{d}t. \end{aligned}$$

Since ψ is strictly convex, with

$$\mathrm{Hess}_x\psi > 2cI,$$

for some positive c and all x in a neighborhood of P, it follows that

$$f_m(x) \geq -\psi(m) + c\|x-m\|^2.$$

Obviously, m is the unique absolute minimum of f_m in P. For the last assertion, we show that the functions

$$\zeta_s := \frac{\mathrm{e}^{-sf_m}}{\|\mathrm{e}^{-sf_m}\|_1}$$

form a Dirac sequence. (We actually show convergence as measures on P.) It is clear from the definition that $\zeta_s > 0$ and $\|\zeta_s\|_1 = 1$, so it remains to show that the norms concentrate around the minimum, that is, given any $\varepsilon, \varepsilon' > 0$ we have to find a s_0 such that

$$\forall s \geq s_0 : \quad \int\limits_{B_\varepsilon(m)} \zeta_s(x)\mathrm{d}x \geq 1 - \varepsilon'.$$

Let $r_0 > 0$ be sufficiently small and let 2α be the maximum of $\mathrm{Hess}_x\psi$ in $B_{r_0}(m)$. Observe that

$$\|\mathrm{e}^{-sf_m}\|_1 = \int_P \mathrm{e}^{-sf_m(x)}\mathrm{d}x \geq \int_{B_r(m)} \mathrm{e}^{-sf_m(x)}\mathrm{d}x \geq d_n r^n \mathrm{e}^{s\psi(m)-s\alpha r^2},$$

for any $r > 0$ such that $r_0 > r$, and where $d_n r^n = \mathrm{Vol}(B_r(m))$. On the other hand,

$$\int\limits_{P\backslash B_\varepsilon(m)} \mathrm{e}^{-sf_m(x)}\mathrm{d}x \leq \int\limits_{P\backslash B_\varepsilon(m)} \mathrm{e}^{s\psi(m)-sc\|x-m\|^2}\mathrm{d}x \leq \mathrm{Vol}(P)\mathrm{e}^{s\psi(m)}\mathrm{e}^{-sc\varepsilon^2}. \tag{19}$$

Therefore,

$$\int\limits_{P\backslash B_\varepsilon(m)} \zeta_s(x)\mathrm{d}x \leq \frac{\mathrm{Vol}(P)\mathrm{e}^{-sc\varepsilon^2+s\alpha r^2}}{d_n r^n}.$$

Choosing r sufficiently small, the right hand side goes to zero as $s \to \infty$ and the result follows. ☐

Let now $\{\sigma_s^m\}_{m\in P\cap\mathbb{Z}^n}$ be the basis of holomorphic sections of L_ω, with respect to the holomorphic structure induced from the map χ_{g_s} in Section 2, that is $\sigma_s^m = \chi_{g_s}^*(\sigma^m)$, for $\sigma^m \in H^0(W_P, L_P)$. We then have

***Proof of Theorem* 1.3.** Using a partition of unity $\{\rho_v\}$ subordinated to the covering by the vertex coordinate charts $\{\check{P}_v\}$, the result can be checked chart by chart. Let $\tau \in C^\infty(L_\omega^{-1})$ be a test section and let

$$\begin{aligned} h_m^s(x) = {}^t(x-m)y - g_s &= \left[{}^t(x-m)(\frac{\partial g_P}{\partial x} + \frac{\partial \varphi}{\partial x}) - g_P - \varphi\right] + s\left[{}^t(x-m)\frac{\partial \psi}{\partial x} - \psi\right] = \\ &= h_m^0(x) + s f_m(x), \end{aligned}$$

with f_m as in Lemma 5.1. Then

$$\begin{aligned} (\iota(\xi_s^m))(\tau) &= \frac{1}{\|\sigma_s^m\|_1}\sum_v \int\limits_{V_v} \rho_v \circ \mu_P(w_v)\mathrm{e}^{-h_m^s\circ\mu_P(w_v)}\mathrm{e}^{\mathrm{i}\ell(m)\vartheta}\tau_v(w_v)\omega^n = \\ &= \frac{1}{\|\sigma_s^m\|_1}\sum_v \int\limits_{\check{P}_v} \rho_v(x)\mathrm{e}^{-h_m^s(x)}\Big(\int\limits_{\mu_P^{-1}(x)} \mathrm{e}^{2\pi\mathrm{i}\ell(m)u}\tau_v(\mathrm{e}^{\frac{\partial g_s}{\partial l}(\ell(x))+2\pi\mathrm{i}u})\mathrm{d}u\Big)\mathrm{d}x = \\ &= \frac{1}{\|\sigma_s^m\|_1}\int\limits_P \mathrm{e}^{-h_m^s(x)}\widehat{\tau}(x,-m)\mathrm{d}x, \end{aligned}$$

where $\widehat{\tau}$ is the fiberwise Fourier transform from equation (17).

Now the L^1-norm in question calculates as

$$\|\sigma_s^m\|_{L^1} = \int\limits_{X_P} \mathrm{e}^{-h_m^s\circ\mu_P}\omega^n = (2\pi)^n\int\limits_P \mathrm{e}^{-h_m^s}\mathrm{d}x.$$

According to Lemma 5.1,

$$\frac{\|\mathrm{e}^{-h_m^0 - sf_m}\|_1}{\|\mathrm{e}^{-sf_m}\|_1} = \int\limits_P \frac{\mathrm{e}^{-sf_m}}{\|\mathrm{e}^{-sf_m}\|_1}\mathrm{e}^{-h_m^0}\mathrm{d}x \to \mathrm{e}^{-h_m^0(m)} \text{ as } s \to \infty,$$

and therefore

$$\iota(\xi_s^m)(\tau) = \int\limits_P \frac{\mathrm{e}^{-h_m^0+sf_m}}{\|\mathrm{e}^{-h_m^0+sf_m}\|_1}\widehat{\tau}(.,-m)\mathrm{d}x \to \widehat{\tau}(m,-m) = \delta^m(\tau)$$

which finishes the proof. ☐

Corollary 5.2. *The results of Lemma 5.1 and Theorem 1.3 are valid for non-compact toric manifolds if one assumes uniform strict convexity of ψ.*

Proof. In Lemma 5.1, the estimate for $f_m(x)$ remains valid for any $x \in P$, so that the function e^{-sf_m} will be integrable even if P is not compact. As for the second part of the proof of Lemma 5.1, instead of (19) one can use

$$\begin{aligned} \int_{P\setminus B_\varepsilon(m)} \mathrm{e}^{-sf_m}\mathrm{d}x &\leq d_n\int_\varepsilon^{+\infty} \mathrm{e}^{s\psi(m)}\mathrm{e}^{-scr^2}r^{n-1}\mathrm{d}r \leq Md_n\int_\varepsilon^{+\infty} \mathrm{e}^{s\psi(m)}\mathrm{e}^{-s\frac{c}{2}r^2}\mathrm{d}r \leq \\ &\leq Md_n\mathrm{e}^{s\psi(m)}\mathrm{e}^{-s\frac{c}{2}\varepsilon^2}\int_0^{+\infty} \mathrm{e}^{-s\frac{c}{2}u^2}\mathrm{d}u \leq M'\mathrm{e}^{s\psi(m)}\mathrm{e}^{-s\frac{c}{2}\varepsilon^2}, \end{aligned}$$

for appropriate constants $M, M' > 0$, where M, M' depend on s but are bounded from above as $s \to \infty$, so that the assertion follows. $\square$

6. COMPACT TROPICAL AMOEBAS

In this section, we undertake a detailed study of the behavior of the compact amoebas in P associated to the family of symplectic potentials in (2)

$$g_s = g_P + \varphi + s\psi,$$

which define the complex structure J_s on X_P, and of their relation to the Log_t amoebas in $\mathbb{R}^n$ [GKZ, Mi, FPT, R].

Let $\breve{Z}_s \subset (\mathbb{C}^*)^n$ be the complex hypersurface defined by the Laurent polynomial

$$\breve{Z}_s = \{w \in (\mathbb{C}^*)^n : \sum_{m \in P \cap \mathbb{Z}^n} a_m \mathrm{e}^{-sv(m)} w^m = 0\},$$

where $a_m \in \mathbb{C}^*, v(m) \in \mathbb{R}$. One natural thing to do in order to obtain the large complex structure limit, consists in introducing the complex structure on $(\mathbb{C}^*)^n$ defined by the complex coordinates $w = e^{z_s}$ where $z_s = sy + i\theta$, and taking the $s \to +\infty$ limit. Then, the map $w \mapsto y$ coincides with the Log_t map for $s = \log t$. However, this deformation of the complex structure, which is well defined for the open dense orbit $(\mathbb{C}^*)^n \subset X_P$ never extends to any (even partial) toric compactification of $(\mathbb{C}^*)^n$. Indeed, that would correspond to rescaling the original symplectic potential by s, which is incompatible with the correct behavior at the boundary of the polytope found by Guillemin and Abreu. Instead, the appropriate thing to do is to consider symplectic potential deformations as above. Even though all complex structures on a compact toric manifold X_P are equivalent, note that by deforming the complex structure while keeping the symplectic structure fixed we are deforming the Kähler metric to, in general, non-equivalent metrics. The $s \to +\infty$-limit shares several properties with the large complex structure limits in the differential geometric description proposed in [KS1, KS2].

As we will describe below, for deformations in the direction of quadratic ψ in (2), in the limit we obtain the Log_t map amoeba intersected with the polytope P. The significative difference is that our limiting tropical amoebas are now compact and live inside P. For more general ψ, they live in the compact image of P by the Legendre transform $\mathcal{L}_\psi$ in (1) and are determined by the locus of non-differentiability of a piecewise linear function, namely as the tropical amoeba of [GKZ, Mi],

$$\mathcal{A}_{\mathrm{trop}} := C^0 - \mathrm{loc}\big(u \mapsto \max_{m \in P \cap \mathbb{Z}^n} \{{}^t m u - v(m)\}\big).$$

6.1. **Limit versus tropical amoebas.**

We are interested in the μ_P-image of the family of (complex) hypersurfaces

$$Y_s := \{p \in X_P : \sum_{m \in P \cap \mathbb{Z}^n} a_m \mathrm{e}^{-sv(m)} \sigma_s^m(p) = 0\} \subset (X_P, J_s)$$

where $a_m \in \mathbb{C}^*$ and $v(m) \in \mathbb{R}$ are parameters and $\sigma_s^m \in H^0\left((X_P, J_s), \chi_{g_s}^* L_P\right)$ is the canonical basis of holomorphic sections of the line bundle $\chi_{g_s}^* L_P$ associated with the polytope P and the symplectic potential (2), introduced in Section 2. We call the image $\mu_P(Y_s) \subset P$ the compact amoeba of Y_s in P.

Definition 6.1. *The limit amoeba* $\mathcal{A}_{\mathrm{lim}}$ *is the subset*

$$\mathcal{A}_{\mathrm{lim}} = \lim_{s\to\infty} \mu_P(Y_s)$$

of the moment polytope P*, where the limit is to be understood as the Hausdorff limit of closed subsets of* P*.*

The existence of this limit is shown in the proof of Theorem 1.7 below. For special large complex structure limits, in particular for $\psi(x) = \frac{x^2}{2}$, we will see that $\mathcal{A}_{\mathrm{lim}}$ itself is piecewise linear. We will relate this amoeba to the tropical (that is, piecewise linear) amoeba of [GKZ, Mi] using a Legendre transform $\check{\chi}_s$ that is the restriction of the map χ_{g_s} described in Section 2 to the open orbit $\check{X}_P$:

$$\begin{tikzcd}
Y_s \arrow[d, phantom, "\cap"] & \check{Y}_s \arrow[l, hook'] \arrow[r, "\cong"] \arrow[d, phantom, "\cap"] & \check{Z}_s \arrow[r, hook] \arrow[d, phantom, "\cap"] & Z_s \arrow[d, phantom, "\cap"] \\
(X_P, \omega, J_s, G_s) \arrow[d, "\mu_P"'] & \check{X}_P \arrow[l, hook'] \arrow[r, "\cong", "\check{\chi}_s"'] \arrow[d, "\mu_P"'] & (\mathbb{C}^*)^n \arrow[r, hook] \arrow[d, "Log_t"] & (W_P, J) \\
P & \check{P} \arrow[l, hook'] \arrow[r, "\cong", "\kappa_s"'] & \mathbb{R}^n &
\end{tikzcd}$$

where κ_s is the family of rescaled Legendre transforms

$$\check{P} \ni x \mapsto \kappa_s(x) := \frac{1}{s}\mathcal{L}_{(g_P+\varphi)+s\psi} = \frac{\partial\psi}{\partial x} + \frac{1}{s}\frac{\partial(g_P+\varphi)}{\partial x} \in \mathbb{R}^n.$$

For any $s > 0$, this is a diffeomorphism $\check{P} \to \mathbb{R}^n$.

Let $\mathcal{A}_s := Log_t(\check{Z}_s)$ be the amoeba of [GKZ, Mi]. Recall that $\mathcal{A}_s \to \mathcal{A}_{\mathrm{trop}}$ in the Hausdorff topology [Mi, R].

Proposition 6.2. *The family of rescaled Legendre transforms* κ_s *satisfies*

$$\kappa_s \circ \mu_P(\check{Y}_s) = \mathcal{A}_s$$

Proof. Under the trivialization of L_P determined by g_s on the open orbit $\check{X}_P$, the sections $\sigma_s^m(x,\theta)$ correspond to polynomial sections w^m, where

$$w = \mathrm{e}^{\frac{\partial(g_P+\varphi)}{\partial x} + s\frac{\partial\psi}{\partial x} + \mathrm{i}\theta}.$$

Combining this with the Log_t-map for $t = \mathrm{e}^s$ gives precisely

$$Log_t w = \frac{\partial\psi}{\partial x} + \frac{1}{s}\frac{\partial(g_P+\varphi)}{\partial x} = \kappa_s(x).$$

□

Remark 6.3. Note that since Y_s is defined as the zero locus of a global section, one has $Y_s = \overline{\check{Y}_s}$ and, in particular, $\mu_P(Y_s) = \overline{\kappa_s^{-1}\mathcal{A}_s}$. On each face F of the moment polytope P, this will consist exactly of the amoeba defined by the sum of monomials corresponding to integer points in F, cf. [Mi].

The family of inverse maps κ_s^{-1} will permit us to capture information not only concerning the open orbit but also the loci of compactification of X_P.

Lemma 6.4. *For any compact subset* $C \subset \check{P}$ *and any* ψ *strictly convex on a neighborhood of* P*,*

$$\kappa_s \to \mathcal{L}_\psi \text{ pointwise on } \check{P} \text{ and uniformly on } C$$

and

$$\kappa_s^{-1} \to \mathcal{L}_\psi^{-1} \textit{ uniformly on } \mathcal{L}_\psi C.$$

Proof. Since $\frac{\partial(g_P+\varphi)}{\partial x}$ is a smooth function on $\breve{P}$, $\kappa_s \to \mathcal{L}_\psi$ pointwise on $\breve{P}$ and uniformly on compact subsets $C \subset \breve{P}$. Furthermore,

$$\|\kappa_s(x) - \kappa_s(x')\| \geq c\|x - x'\|, \qquad \forall x, x' \in \breve{P}, \tag{20}$$

with a constant $c > 0$ uniform in s, since the derivative

$$\frac{\partial \kappa_s}{\partial x} = \text{Hess}_x \psi + \frac{1}{s}\text{Hess}_x(g_P + \varphi) > \text{Hess}_x \psi > 0$$

is (uniformly) positive definite. Therefore, the family of inverse mappings κ_s^{-1} is uniformly Lipschitz (on $\mathbb{R}^n$). Thus the pointwise convergence $\kappa_s^{-1} \to \mathcal{L}_\psi^{-1}$ on $\mathcal{L}_\psi \breve{P}$ is uniform on any compact $\mathcal{L}_\psi C$. $\square$

Before proving the main theorem, we recall some facts about convex sets in $\mathbb{R}^n$ and also show two auxiliary lemmata on the behavior of the gradient of any toric symplectic potential near the boundary of the moment polytope. Consider any constant metric $G = {}^tG > 0$ on $\mathbb{R}^n$. For an arbitrary closed convex polyhedral set $P \subset \mathbb{R}^n$ and any point $p \in \partial P$, denote by $\mathcal{C}_p^G$ the closed cone of directions that are "outward pointing at p" in the following sense,

$$\mathcal{C}_p^G(P) := \{c \in \mathbb{R}^n : {}^tcG(p - p') \geq 0, \quad \forall p' \in P\}.$$

Notice that for the Euclidean metric $G = I$, the cone of p is precisely the negative of the cone of the fan of P, corresponding to the face p lies in; in this case we will write $\mathcal{C}_p := \mathcal{C}_p^I$.

Lemma 6.5. *For any sequence $x_k \in \breve{P}$ that converges to a point in the boundary, $x_k \to p \in \partial P$, we have*

$$\frac{\partial(g_P + \varphi)}{\partial x}|_{x_k} \to \mathcal{C}_p(P),$$

in the sense that for any $c \notin \mathcal{C}_p(P)$ there is an open neighborhood $U \ni p$ such that

$$\mathbb{R}_0^+ \frac{\partial(g_P + \varphi)}{\partial x}|_x \neq \mathbb{R}_0^+ c \qquad \forall x \in U \cap \breve{P}.$$

Proof. Suppose (using an affine change of coordinates $l(x) = Ax - \lambda$) that p lies in the codimension k face, $k > 0$, where $l_1 = \cdots = l_k = 0$ and $l_j > 0$ for $j = k+1, \ldots, n$. We have

$$\mathcal{C}_p = \{c \in \mathbb{R}^n : {}^tcA^{-1}(l(p) - l(p')) \geq 0, \forall p' \in P\},$$

and $l_i(p - p') \leq 0$ for $i = 1, \ldots, k$, whereas there is no restriction on the sign of $l_j(p - p')$ for $j = k+1, \ldots, n$. Therefore,

$$c \in \mathcal{C}_p \iff c = {}^tA\widetilde{c} \text{ with } \widetilde{c} \in (\mathbb{R}_0^-)^k \times \{0\} \subset \mathbb{R}^n.$$

Since

$$\frac{\partial(g_P + \varphi)}{\partial x} = {}^tA\frac{\partial(g_P + \varphi)}{\partial l}$$

it is therefore sufficent to prove that $\frac{\partial(g_P+\varphi)}{\partial l}$ approaches $(\mathbb{R}_0^-)^k \times \{0\} \subset \mathbb{R}^n$ as we get near p. Indeed, we find that

$$\frac{\partial g_P}{\partial l} = \frac{1}{2}\frac{\partial}{\partial l}\sum_{a=1}^{d} \ell_a \log \ell_a = \frac{1}{2}\frac{\partial}{\partial l}\Big(\sum_{i=1}^{k} l_i \log l_i + \sum_{j=k+1}^{n} l_j \log l_j + \sum_{m=n+1}^{d} \ell_m \log \ell_m\Big)$$

and hence

$$\frac{\partial g_P}{\partial l_r} = \frac{1}{2}(1 + \log l_r + \sum_{m=n+1}^{d} \frac{\partial \ell_m}{\partial l_r}(1+\log \ell_m)). \tag{21}$$

For $m > n$ (actually, for $m > k$), the sum is bounded in a neighborhood of p since $\frac{\partial \ell_m}{\partial l_r}$ is constant and $\ell_m > 0$ at p. Since $\frac{\partial \varphi}{\partial l_r}$ is also bounded for any r, $\frac{\partial(g_P+\varphi)}{\partial l_j}$ is bounded in a neighborhood of p for $j = k+1, \ldots, n$ and clearly $\frac{\partial(g_P+\varphi)}{\partial l_i} \to -\infty$ for $i = 1, \ldots, k$ as we approach p, which proves the lemma. ☐

In the following lemma, we relate the Legendre transforms κ_s and $\mathcal{L}_\psi$ at large s, more precisely:

Lemma 6.6. *For any two points $p \neq p' \in P$, there exist $\varepsilon > 0$ and $s_0 \geq 0$ that depend only on p' and the distance $d(p,p')$, such that for all $s \geq s_0$*

$$\overline{\kappa_s(B_\varepsilon(p') \cap \breve{P})} \cap \big(\mathcal{L}_\psi(p) + \mathcal{C}_p(P)\big) = \emptyset.$$

Proof. We will show that there is a hyperplane separating $\kappa_s(B_\varepsilon(p') \cap \breve{P})$ and $\mathcal{L}_\psi(p) + \mathcal{C}_p(P)$; we continue to use a chart as in Lemma 6.5.

For any two points $p \neq p'$,

$$\begin{aligned} {}^t(l(p)-l(p'))(\frac{\partial \psi}{\partial l}|_p - \frac{\partial \psi}{\partial l}|_{p'}) &= {}^t(l(p)-l(p'))\int_0^1 (\mathrm{Hess}_{\,p'+\tau(p-p')}\psi)\tau \mathrm{d}\tau(l(p)-l(p')) \geq \\ &\geq c_\psi \|l(p)-l(p')\|^2 > 0 \end{aligned}$$

where $c_\psi > 0$ is a constant depending only ψ. This implies that there is at least one index j such that $l_j(p) \neq l_j(p')$,

$$\mathrm{sign}(\frac{\partial \psi}{\partial l_j}|_p - \frac{\partial \psi}{\partial l_j}|_{p'}) = \mathrm{sign}(l_j(p) - l_j(p')),$$

and also

$$\left|\frac{\partial \psi}{\partial l_j}|_p - \frac{\partial \psi}{\partial l_j}|_{p'}\right| \geq \frac{c_\psi}{n}|l_j(p) - l_j(p')|.$$

Choose $\varepsilon > 0$ small enough (this choice depends on ψ only) so that

$$\frac{\partial \psi}{\partial l_j}|_p \notin \frac{\partial \psi}{\partial l_j}(B_\varepsilon(p') \cap \breve{P})$$

and consider first the case that $0 \leq l_j(p) < l_j(p')$. For all $x \in B_\varepsilon(p') \cap \breve{P}$, from equation (21),

$$\frac{\partial g_P}{\partial l_j}|_x \geq c_1 \log^-(l_j(p') - \varepsilon) + c_2 = C,$$

where $\log^-$ denotes the negative part of the logarithm, and c_1, c_2, C are constants depending only on p' and ε. Then, for any $\delta > 0$ such that

$$\frac{\partial \psi}{\partial l_j}|_p + \delta \notin \frac{\partial \psi}{\partial l_j}(B_\varepsilon(p') \cap \breve{P}),$$

we find $s_0 = \frac{2|C|}{\delta}$ so that

$$\frac{\partial\psi}{\partial l_j}|_x + \frac{1}{s}\frac{\partial g_P}{\partial l_j}|_x \geq \frac{\partial\psi}{\partial l_j}(p) + \frac{\delta}{2}, \qquad \forall x \in B_\varepsilon(p') \cap \check{P}.$$

Hence, also

$$\frac{\partial(\psi + \frac{1}{s}\varphi)}{\partial l_j}|_x + \frac{1}{s}\frac{\partial g_P}{\partial l_j}|_x \geq \frac{\partial\psi}{\partial l_j}(p) + \frac{\delta}{4}, \qquad \forall x \in B_\varepsilon(p') \cap \check{P},$$

for s big enough (the additional condition depending only on φ, which is globally controlled on the whole polytope P), which proves our assertion.

If, on the other hand, $0 \leq l_j(p') < l_j(p)$, we see again from equation (21) that

$$\frac{\partial g_P}{\partial l_j} \leq c_1' \log^+ l_j + c_2'$$

on $B_\varepsilon(p') \cap \check{P}$, where $\log^+$ stands for the positive part of the logarithm. Again,

$$\frac{\partial g_P}{\partial l_j}|_x \leq c_1' \log^+(l_j(p') + \varepsilon) + c_2' = C', \qquad \forall x \in B_\varepsilon(p') \cap \check{P}$$

and the same argument applies. □

We will now characterize the limit amoeba in terms of the tropical amoeba via a projection π that can be described as follows.

Lemma 6.7. *For any strictly convex ψ as above, there exists a partition of $\mathbb{R}^n$ indexed by P of the form*

$$\mathbb{R}^n = \coprod_{p\in P} \mathcal{L}_\psi(p) + \mathcal{C}_p(P). \tag{22}$$

In particular, there is a well-defined continuous projection $\pi : \mathbb{R}^n \to \mathcal{L}_\psi P$ given by

$$\pi(\mathcal{L}_\psi(p) + \mathcal{C}_p(P)) = \mathcal{L}_\psi(p).$$

Proof. It suffices to show that for $p \neq p'$,

$$\big(\mathcal{L}_\psi(p) + \mathcal{C}_p(P)\big) \cap \big(\mathcal{L}_\psi(p') + \mathcal{C}_{p'}(P)\big) = \emptyset.$$

To see this, assume that

$$\mathcal{L}_\psi(p) + c = \mathcal{L}_\psi(p') + c', \text{ with } c \in \mathcal{C}_p(P), c' \in \mathcal{C}_{p'}(P).$$

Then ${}^t(c - c')(p - p') \geq 0$, from the definition of the cones; on the other hand,

$$\begin{aligned} {}^t(p-p')(c-c') &= {}^t(p-p')(\mathcal{L}_\psi(p') - \mathcal{L}_\psi(p)) = \\ &= {}^t(p-p')\int_0^1 (\mathrm{Hess}_{\,p+\tau(p'-p)}\psi)\mathrm{d}\tau(p'-p) < 0, \end{aligned}$$

which is a contradiction. □

Remark 6.8. For quadratic ψ with ${}^tG = G > 0$ symmetric and positive definite there is a more intrinsic description of the map π: it is given by the projection of $\mathbb{R}^n$ on the closed convex subset P under which each point projects onto its best approximation in the polytope $\mathcal{L}_\psi P$ with respect to the metric G^{-1} (see, for instance, chapter v of [Bou], and also Figure 1),

$$p = \pi(y) \iff p \in \mathcal{L}_\psi P \wedge \forall p' \in \mathcal{L}_\psi P \setminus \{p\} : \|y - p\|_{G^{-1}} < \|y - p'\|_{G^{-1}}.$$



FIGURE 1. The map π.

Note also that

$$\forall y \in \mathbb{R}^n : y - \pi(y) \in \mathcal{C}_{\pi(y)}^{G^{-1}}(\mathcal{L}_\psi P)$$

and that, in fact, $\pi(y)$ is characterised by this property, i.e.

$$\forall p \in \mathcal{L}_\psi P : y - p \in \mathcal{C}_p^{G^{-1}}(\mathcal{L}_\psi P) \iff \pi(y) = p.$$

In this sense, $\mathcal{C}_p^{G^{-1}}(\mathcal{L}_\psi P)$ is a kind of "convex kernel at p" of the convex projection π. Note, by the way, that in this case $\mathcal{C}_x(P) = \mathcal{C}_{\mathcal{L}_\psi(x)}^{G^{-1}}(\mathcal{L}_\psi P)$.

Finally, we have

***Proof of Theorem* 1.7.** We first show that $\mathcal{L}_\psi \circ \kappa_s^{-1} \to \pi$ pointwise on $\mathbb{R}^n$. For points in the interior of $\mathcal{L}_\psi P$, where $\pi|_{\breve{P}} = \mathrm{id}_{\breve{P}}$, this is clear from Lemma 6.4.

Consider, therefore, any point $y \notin \mathcal{L}_\psi \breve{P}$, its family of inverse images $x_s = \kappa_s^{-1}(y) \in \breve{P}$, and any convergent subsequence $x_{s_k} \to p$. Then the limit lies in the boundary, $p \in \partial P$. We need to show that $\mathcal{L}_\psi(p) = \pi(y)$, or, what is the same, that $y - \mathcal{L}_\psi(p) \in \mathcal{C}_p(P)$. This is guaranteed by Lemma 6.5,

$$\frac{1}{s}\frac{\partial(g_P + \varphi)}{\partial x}|_{x_{s_k}} = (\kappa_s - \mathcal{L}_\psi)(x_{s_k}) = y - \mathcal{L}_\psi(x_{s_k}) \to y - \mathcal{L}_\psi(p) \in \mathcal{C}_p(P),$$

and proves pointwise convergence.

Now we can use compactness of P (and hence of the space of closed non-empty subsets of P with the Hausdorff metric) to show the result. Throughout the proof we will not distinguish between sets and their closures since the Hausdorff topology does not separate them.

Let us first show that

$$\kappa_s^{-1}\mathcal{A}_{\mathrm{trop}} \xrightarrow{\mathrm{H}} \mathcal{L}_\psi^{-1} \circ \pi \mathcal{A}_{\mathrm{trop}} \quad .$$

Take any convergent subsequence $\kappa_{s_k}^{-1}\mathcal{A}_{\mathrm{trop}} \xrightarrow{\mathrm{H}} K \subset P$; since $\kappa_s^{-1} \to \mathcal{L}_\psi^{-1} \circ \pi$ pointwise, it follows that

$$K \supset \mathcal{L}_\psi^{-1} \circ \pi \mathcal{A}_{\mathrm{trop}}.$$

For the other inclusion, consider any point $p' \notin \mathcal{L}_\psi^{-1} \circ \pi \mathcal{A}_{\mathrm{trop}}$; since the distance of p' to $\mathcal{L}_\psi^{-1} \circ \pi \mathcal{A}_{\mathrm{trop}}$ is strictly positive, by Lemma 6.6 there is a neighborhood U of p' in $\breve{P}$ and a s_0 such that for all $p \in \mathcal{L}_\psi^{-1} \circ \pi\mathcal{A}_{\text{trop}}$ and $s \geq s_0$, the sets $\kappa_s(U)$ and $\mathcal{L}_\psi(p) + \mathcal{C}_p(P)$ not only have empty intersection but are actually separated by a hyperplane. But this implies, in particular, that for s large enough

$$U \cap \kappa_s^{-1}\mathcal{A}_{\text{trop}} = \emptyset$$

and $p' \notin K$, as we wished to show.

In the last step, we prove that $\kappa_s^{-1}\mathcal{A}_s \xrightarrow{\text{H}} \mathcal{L}_\psi^{-1} \circ \pi\mathcal{A}_{\text{trop}}$. Again, using compactness, it is sufficient to consider any convergent subsequence $\kappa_{s_k}^{-1}\mathcal{A}_{s_k} \xrightarrow{\text{H}} K'$.

To show that $\mathcal{L}_\psi^{-1} \circ \pi\mathcal{A}_{\text{trop}} \subset K'$, it is sufficent to observe that $\mathcal{A}_{\text{trop}} \subset \mathcal{A}_{s_k}$ (see [GKZ, Mi]) and hence $\kappa_{s_k}^{-1}\mathcal{A}_{\text{trop}} \subset \kappa_{s_k}^{-1}\mathcal{A}_{s_k}$.

For the converse inclusion $K' \subset \mathcal{L}_\psi^{-1} \circ \pi\mathcal{A}_{\text{trop}}$, consider the constant c from inequality (20) above, and set

$$\varepsilon_k := \frac{1}{c}\text{dist}(\mathcal{A}_{s_k}, \mathcal{A}_{\text{trop}}).$$

This sequence converges to zero, and therefore the closed ε_k-neighborhoods $(\kappa_{s_k}^{-1}\mathcal{A}_{\text{trop}})_{\varepsilon_k} \supset \kappa_{s_k}^{-1}\mathcal{A}_{\text{trop}}$ still converge to $\mathcal{L}_\psi^{-1} \circ \pi\mathcal{A}_{\text{trop}}$. But as κ_{s_k} satisfies the uniform bound in (20),

$$\kappa_{s_k}((\kappa_{s_k}^{-1}\mathcal{A}_{\text{trop}})_{\varepsilon_k}) \supset (\mathcal{A}_{\text{trop}})_{c\varepsilon_k} \supset \mathcal{A}_{s_k}$$

and hence

$$(\kappa_{s_k}^{-1}\mathcal{A}_{\text{trop}})_{\varepsilon_k} \supset \kappa_{s_k}^{-1}\mathcal{A}_{s_k}$$

which proves the second inclusion. □

Remark 6.9. Note that for quadratic ψ, the limit amoeba $\mathcal{A}_{\text{lim}} \subset P$ itself is piecewise linear.

Remark 6.10. Note that there is an open set of valuations v and directions ψ such that the projection $\pi\mathcal{A}_{\text{trop}}$ will coincide with $\mathcal{L}_\psi P \cap \mathcal{A}_{\text{trop}}$; this happens whenever the "nucleus" of the tropical amoeba $\mathcal{A}_{\text{trop}}$ lies inside $L_\psi P$, since the infinite legs run off to infinity along directions in the cone of the relevant faces. This situation is depicted in Figure 2 for a limit of elliptic curves in a del Pezzo surface,

$$\begin{aligned} P &= \text{conv}\{(1,0),(1,1),(0,1),(-1,0),(-1,-1),(0,-1)\} \\ \psi(x) &= \frac{x^2}{2} + \frac{\|x\|^4}{4} \\ v(m) &= \frac{m^2}{2}. \end{aligned}$$

Remark 6.11. The behavior of the limit of $\mu_P(Y_s)$ will therefore vary greatly with ψ, even with the valuation $v(m)$ held fixed. We illustrate this with the simplest possible example, $\mathbb{P}^2$, with moment polytope the standard simplex in $\mathbb{R}^2$ and valuation $v(0,0) = 0$, $v(1,0) = \frac{1}{2}$, $v(0,1) = \frac{1}{4}$, in Figure 3.

6.2. **Implosion of polytopes versus explosion of fans.** For simplicity, in the present subsection, we restrict ourselves to the case $\psi(x) = \frac{1}{2}x^2$. We remark that while the map π projects onto P, the map $id - \pi$ is injective in the interior of the cones $v + \mathcal{C}_v$ for all vertices $v \in P$ (regions $1-4$ in figure 4). For a face F of dimension $k > 0$ of P, the region $\breve{F} + \mathcal{C}_p$, for any $p \in \breve{F}$, implodes to the cone $\mathcal{C}_p$ of codimension k. In particular,



Figure 2. The situation of the main theorem (for ψ not quadratic): $P \supset \mathcal{A}_{\mathrm{lim}}$ (top left), $\mathcal{L}_\psi P$ with the cones of projection $\mathcal{C}_p$ (dotted) and the tropical amoeba $\mathcal{A}_{\mathrm{trop}}$ (top right) and $\mathcal{L}_\psi P \supset \pi\mathcal{A}_{\mathrm{trop}} = \mathcal{L}_\psi\mathcal{A}_{\mathrm{lim}}$ (bottom).

the polytope P implodes to the origin. Dually, the map $id - \pi$ explodes the fan along positive codimension cones. In particular, the origin is exploded to P. In Figure 4, we consider the non-generic Laurent polynomial

$$a_1 + a_2 t^{-0.6} x + a_3 t^{-0.4} x^2 + a_4 t^{1.8} \frac{y^2}{x},$$

its tropical amoeba for $\mathbb{P}^2$ blown-up at one point and the corresponding images under the maps π and $id - \pi$. Note that, for this non generic polynomial, part of the compact amoeba $\pi\mathcal{A}_{\mathrm{trop}}$ lies in the boundary of P. Only in $\check{P}$ does $\pi\mathcal{A}_{\mathrm{trop}}$ coincide with the tropical non-Archimedean amoeba.

6.3. **Amoebas associated to geometric quantization.** As we could observe in Example 6.11, the behavior of the limit amoeba defined via a fixed valuation is rather unstable. Actually, it does not only depend on the choice of ψ, but also behaves badly under integer translations of the moment polytope P. These shortcomings are somehow overcome by a specific choice of valuation that is associated with a toric variety and a large complex structure limit with quadratic ψ. This construction provides the natural link between the convergence of sections to delta distributions considered in Section 5 that comes out of geometric quantization, and the consideration of the image of hypersurfaces defined by the zero locus of generic sections.

Geometric quantization motivates considering the hypersurfaces defined by

$$\check{Y}_s^{\mathrm{GQ}} = \{p \in X_P : \sum_{m \in P \cap \mathbb{Z}^n} a_m \xi_s^m(p) = 0\},$$

where $a_m \in \mathbb{C}^*$, and ξ_s^m are the L^1-normalized holomorphic sections converging to delta distributions, as in Section 5. A simple estimate of the order of decay of $\|\sigma_s^m\|_1$ as


$\mathcal{L}_\psi = \mathrm{id}$
$\mathcal{L}_\psi = \frac{1}{4}\begin{bmatrix} 6 & 3 \\ 3 & 6 \end{bmatrix}$
$\mathcal{L}_\psi = \frac{1}{4}\begin{bmatrix} 3 & 2 \\ 2 & 3 \end{bmatrix}$
π
π
π

Figure 3. Limit amoebas ($\mathcal{A}_{\mathrm{lim}} \subset P$, top row), their image under $\mathcal{L}_\psi$, and tropical amoebas ($\mathcal{A}_{\mathrm{trop}}$, bottom row) for different quadratic ψ and fixed valuation v.

$s \to \infty$ gives, in the notation of Lemma 5.1,

$$\begin{aligned} \|\sigma_s^m\|_1^{-1}\frac{\mathrm{d}}{\mathrm{d}s}\|\sigma_s^m\|_1 &= \|\sigma_s^m\|_1^{-1}\frac{\mathrm{d}}{\mathrm{d}s}\int\limits_P \mathrm{e}^{-h_m^s(x)}\mathrm{d}x = \\ &= \|\mathrm{e}^{-h_m^s}\|_1^{-1}\frac{\mathrm{d}}{\mathrm{d}s}\int\limits_P \mathrm{e}^{-h_m^0(x)-sf_m(x)}\mathrm{d}x = \\ &= -\int\limits_P f_m(x)\frac{\mathrm{e}^{-h_m^s(x)}}{\|\mathrm{e}^{-h_m^s}\|_1}\mathrm{d}x \to -f_m(m) = \psi(m). \end{aligned}$$

Therefore, we call the limit of the family of amoebas $\mu_P(Y_s)$, where

$$\check{Y}_s = \{w \in (\mathbb{C}^*)^n : \sum_{m \in P\cap\mathbb{Z}^n} a_m \mathrm{e}^{-s\psi(m)} w^m = 0\},$$

the GQ limit amoeba, $\mathcal{A}_{\mathrm{lim}}^{\mathrm{GQ}}$. The fact that for this choice of valuations, inspired by geometric quantization, the limit amoeba keeps away from integral points in P is consistent with the convergence of the holomorphic sections ξ_s^m to delta distributions supported on the Bohr-Sommerfeld fibers corresponding to those integral points.

The behavior of this "natural" (from the point of view of geometric quantization) choice of valuation is illustrated in Figure 5 for different quadratic ψ. Note, in particular, that in the case G_2 there are parts of the tropical amoeba $\mathcal{A}_{\mathrm{trop}}$ that lie outside $\mathcal{L}_\psi P$ and



Figure 4. Explosion of the fan and implosion of the polytope. The marked points correspond to non-zero coefficients a_m.

get projected onto subsets of faces (with non-empty interior in the relative topology). Below, we will give a complete characterization of the GQ limit amoeba in this situation.



$G_0 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$ $G_1 = \frac{1}{4}\begin{bmatrix} 6 & 3 \\ 3 & 6 \end{bmatrix}$ $G_2 = G_1^{-1} = \frac{4}{9}\begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$

Figure 5. GQ amoebas associated to different quadratic ψ's.

We start by observing that from the point of view of equivalence of toric varietes described by different Delzant polytopes, the GQ limit amoeba is well behaved:

Proposition 6.12. *Let* $\psi \in C^\infty_+(P)$.

a. *For any integer vector* $k \in \mathbb{Z}^n$*, setting* $\widetilde{P} = P + k$ *and* $\widetilde{\psi}(\widetilde{x}) = \psi(\widetilde{x} - k)$*, we have*

$$\widetilde{\mathcal{A}}^{\mathrm{GQ}}_{\mathrm{lim}} = \mathcal{A}^{\mathrm{GQ}}_{\mathrm{lim}} + k,$$

b. *For any base change* $A \in Sl(n, \mathbb{Z})$ *of the lattice* $\mathbb{Z}^n$*, setting* $\widetilde{P} = AP$ *and* $\widetilde{\psi}(\widetilde{x}) = \psi(A^{-1}\widetilde{x})$*, we have*

$$\widetilde{\mathcal{A}}^{\mathrm{GQ}}_{\mathrm{lim}} = A\mathcal{A}^{\mathrm{GQ}}_{\mathrm{lim}}.$$

Proof. a.: since $\widetilde{\psi}(\widetilde{m} = m + k) = \psi(m)$, $\widetilde{\mathcal{A}}_{\mathrm{trop}} = \mathcal{A}_{\mathrm{trop}}$; on the other hand,

$$\frac{\partial \widetilde{\psi}}{\partial x}|_{\widetilde{x}=x+k} = \frac{\partial \psi}{\partial x}|_x$$

thus $\mathcal{L}_\psi P = \mathcal{L}_{\widetilde{\psi}} \widetilde{P}$ and

$$\mathcal{L}_\psi P \cap \mathcal{A}_{\mathrm{trop}} = \mathcal{L}_{\widetilde{\psi}} \widetilde{P} \cap \widetilde{\mathcal{A}}_{\mathrm{trop}}.$$

b.: similarly, since $\widetilde{\psi}(\widetilde{m} = Am) = \psi(m)$ and the tropical amoeba $\widetilde{\mathcal{A}}_{\mathrm{trop}}$ is defined via the functions

$$\widetilde{u} \mapsto {}^t\widetilde{m}\widetilde{u} - \widetilde{\psi}(\widetilde{m}) = {}^tm{}^tA\widetilde{u} - \psi(m)$$

it follows that $\widetilde{\mathcal{A}}_{\mathrm{trop}} = {}^tA^{-1}\mathcal{A}_{\mathrm{trop}}$; for the Legendre transforms one finds

$$\frac{\partial \widetilde{\psi}}{\partial x}|_{\widetilde{x}=Ax} = {}^tA^{-1}\frac{\partial \psi}{\partial x}|_x$$

which proves the second claim. □

Actually, as is to be expected from the convergence of the sections defining $\mathcal{A}^{\mathrm{GQ}}_{\mathrm{lim}}$ to delta distributions, these amoebas never intersect lattice points:

Proposition 6.13. *For any strictly convex* $\psi \in C^\infty_+(P)$*, the GQ amoeba* $\mathcal{A}^{\mathrm{GQ}}_{\mathrm{lim}}$ *stays away from lattice points in the interior of* P*, that is,*

$$\mathcal{A}^{\mathrm{GQ}}_{\mathrm{lim}} \cap \breve{P} \cap \mathbb{Z}^n = \emptyset.$$

Proof. We consider the functions that were used to define the tropical amoeba,

$$\eta_m(u) = {}^tmu - \psi(m), \qquad \forall m \in P \cap \mathbb{Z}^n$$

and observe that for $u = u_m = \mathcal{L}_\psi m = \frac{\partial \psi}{\partial x}|_m$ this is the value of the Legendre transform h at u_m,

$$\eta_m(u_m) = h(u_m)$$

where $h(u) = {}^tx(u)u - \psi(x(u)$. To show that

$$\eta_m(u_m) > \eta_{\widetilde{m}}(u_m) \qquad \forall \widetilde{m} \in P \cap \mathbb{Z}^n,\ \widetilde{m} \neq m$$

we use convexity of the Legendre transform $u \mapsto h(u)$, namely

$$\begin{aligned} \eta_m(u_m) = h(u_m) \quad &> \quad h(u_{\widetilde{m}}) + {}^t(u_m - u_{\widetilde{m}})\frac{\partial h}{\partial u}|_{u_{\widetilde{m}}} = \\ &= \quad {}^t\widetilde{m}u_{\widetilde{m}} - \psi(\widetilde{m}) + {}^t(u_m - u_{\widetilde{m}})\widetilde{m} = \eta_{\widetilde{m}}(u_m) \end{aligned}$$

where we used the fact that

$$\frac{\partial h}{\partial u}|_{u_{\widetilde{m}}} = \frac{\partial h}{\partial u}|_{\frac{\partial \psi}{\partial x}|_{\widetilde{m}}} = \widetilde{m}.$$

□

When ψ is quadratic it is possible to characterize the GQ limit amoeba completely in terms of the limit metric on P only. Let F_p denote the minimal face containing any given point $p \in \partial P$. Note that for any point $x \in P$,

$$x \in F_p \iff (x-p) \perp_G \mathcal{C}_p^G$$

and, more generally, for any $x \in P$ and $c \in \mathcal{C}_p^G$

$$\|x+c-p\|_G^2 = \|x-p\|_G^2 + \|c\|_G^2 - 2\|x-p\|_G\|c\|_G \cos\alpha \tag{23}$$

where $\alpha = \angle_G(x-p,c) \geq \frac{\pi}{2}$ since ${}^t cG(p-x) \geq 0$ by definition of $\mathcal{C}_p^G(P)$.

Proposition 6.14. *Let $\psi(x) = \frac{{}^t xGx}{2} + {}^t bx$ with ${}^t G = G > 0$; then a point $p \in P$ belongs to $\mathcal{A}_{\mathrm{lim}}^{\mathrm{GQ}}$ if and only if one of the following conditions holds:*

- **a.** *There are (at least) two lattice points $m_1 \neq m_2 \in P \cap \mathbb{Z}^n$ such that*
$$\|p-m_1\|_G = \|p-m_2\|_G = \min_{m \in P \cap \mathbb{Z}^n}\{\|p-m\|_G\}.$$
- **b.** *$p \in \partial P$ and the unique closest lattice point does not lie in the face F_p.*

Remark 6.15. The two conditions are, evidently, mutually exclusive: from the description of the map π it is evident that the inverse image of the intersection of the tropical amoeba $\mathcal{A}_{\mathrm{trop}}$ with $\mathcal{L}_\psi P$ is always a subset of $\mathcal{A}_{\mathrm{lim}}$. This is taken care of by a., while b. describes the parts of the GQ limit amoeba that arise from parts of $\mathcal{A}_{\mathrm{trop}}$ that are "smashed on the boundary" by the convex projection. In particular, for points in the interior condition b. is irrelevant.

Proof. a.: For quadratic ψ and $u = \mathcal{L}_\psi x = Gx + b$, take

$$\begin{aligned} \eta_m(u) &= {}^t mu - \psi(m) = {}^t mu - (\frac{{}^t mGm}{2} + {}^t bm) = \\ &= \big({}^t(\mathcal{L}_\psi m)G^{-1}u - {}^t bG^{-1}u\big) - \frac{1}{2}\big(\|\mathcal{L}_\psi m\|_{G^{-1}}^2 - \|b\|_{G^{-1}}^2\big) = \\ &= \frac{1}{2}\big(\|u-b\|_{G^{-1}}^2 - \|u-\mathcal{L}_\psi m\|_{G^{-1}}^2\big). \end{aligned}$$

Since the first term is independent of m, it is irrelevant for the locus of non-differentiability that defines the tropical amoeba,

$$\begin{aligned} \mathcal{A}_{\mathrm{trop}} &= C^0-\mathrm{loc}\big(u \mapsto \max_{m \in P\cap\mathbb{Z}^n}\{\eta_m(u)\}\big) = \\ &= C^0-\mathrm{loc}\big(u \mapsto \frac{1}{2}\|u-b\|_{G^{-1}}^2 - \min_{m\in P\cap\mathbb{Z}^n}\{\frac{1}{2}\|u-\mathcal{L}_\psi m\|_{G^{-1}}^2\}\big). \end{aligned}$$

Therefore, u lies in the tropical amoeba if and only if there are two distinct lattice points $m_1 \neq m_2$ in P such that

$$\|u-\mathcal{L}_\psi m_1\|_{G^{-1}} = \|u-\mathcal{L}_\psi m_2\|_{G^{-1}} \iff \|p-m_1\|_G = \|p-m_2\|_G$$

for $u = \mathcal{L}_\psi p \in \mathcal{L}_\psi P$. Taking into account that $\mathcal{L}_\psi \mathcal{A}_{\mathrm{lim}} \supset \mathcal{A}_{\mathrm{trop}} \cap \mathcal{L}_\psi P$, this proves that condition a. is necessary and sufficient for u to belong to this intersection. Thus, either p satisfies a. or if it belongs to $\mathcal{A}_{\mathrm{lim}}$ then it belongs to $\mathcal{A}_{\mathrm{lim}} \setminus \mathcal{A}_{\mathrm{trop}}$.

b.: Fix $p \in \partial P$. First we show that if there is a unique nearest lattice point, say m_p, that lies in the face of p, F_p, then $p \notin \mathcal{A}_{\mathrm{lim}}$. By the definition of $\mathcal{A}_{\mathrm{lim}}$, we have to show that

$$\forall c \in \mathcal{C}_p^G(P): \qquad \mathcal{L}_\psi p + c \notin \mathcal{A}_{\mathrm{trop}},$$

which follows in particular if $\eta_{m_p}(\mathcal{L}_\psi p+c) > \eta_m(\mathcal{L}_\psi p+c)$ for any lattice point $m \neq m_p$. By the reasoning above, this is equivalent to

$$\|\mathcal{L}_\psi p + c - \mathcal{L}_\psi m_p\|^2_{G^{-1}} < \|\mathcal{L}_\psi p + c - \mathcal{L}_\psi m\|^2_{G^{-1}},$$

which follows straight from equation (23) (with $\cos\alpha = 0$).

For the other implication, assume $m_p \notin F_p$. Then,

$$\eta_{m_p}(\mathcal{L}_\psi p) > \eta_m(\mathcal{L}_\psi p), \qquad \forall m \in P \cap \mathbb{Z}^n, m \neq m_p$$

and it suffices to show that for any $m \in F_p \cap \mathbb{Z}^n$ and $c \in \mathcal{C}^G_p(P) \setminus \{0\}$

$$\|\mathcal{L}_\psi p + \tau c - \mathcal{L}_\psi m\|^2_{G^{-1}} < \|\mathcal{L}_\psi p + \tau c - \mathcal{L}_\psi m_p\|^2_{G^{-1}} \tag{24}$$

for some $\tau > 0$ large enough. The left hand side equals

$$\|\mathcal{L}_\psi p + \tau c - \mathcal{L}_\psi m\|^2_{G^{-1}} = \|p-m\|^2_G + \tau^2\|c\|^2_{G^{-1}}$$

whereas the right hand side gives

$$\|\mathcal{L}_\psi p + \tau c - \mathcal{L}_\psi m_p\|^2_{G^{-1}} = \|p-m_p\|^2_G + \tau^2\|c\|^2_{G^{-1}} - 2\tau\|p-m_p\|_G\|c\|_{G^{-1}}\cos\alpha$$

where $\cos\alpha < 0$. Subtracting therefore the left hand side of inequality (24) from the right hand side, we are left with the expression

$$\|\mathcal{L}_\psi p + \tau c - \mathcal{L}_\psi m_p\|^2_{G^{-1}} - \|\mathcal{L}_\psi p + \tau c - \mathcal{L}_\psi m\|^2_{G^{-1}} = -2\tau\|p-m_p\|_G\|c\|_{G^{-1}}\cos\alpha \to +\infty,$$

as $\tau \to \infty$, which finishes the proof. □

6.4. Relation to other aspects of degeneration of complex structures.

Degenerating families of complex structures have been studied from a variety of viewpoints. In this section we would like to briefly address aspects of the relation of the present work to some of those.

The first link, is to the occurance of torus fibrations in mirror symmetry, following [SYZ]. For γ_s as in (9), with

$$G_s = \mathrm{Hess}\,(g_P + \varphi + s\psi),$$

consider the rescaled metric

$$\eta_s = \frac{1}{\mathrm{diam}^2(X_P, \gamma_s)}\gamma_s.$$

Then, as $s \to \infty$, in the metric η_s the fibers of the moment map μ_P collapse to the base P. Note that, as $s \to \infty$, η_s gives a well defined metric on $\breve{P}$,

$$G_{\lim} = \frac{1}{b}\mathrm{Hess}\,\psi,$$

where $b = \lim_{s\to\infty}\frac{\mathrm{diam}^2(X_P,\gamma_s)}{s}$. Let now $i : \mathcal{A}_{\lim} \hookrightarrow P$ be the embedding of $\mathcal{A}_{\lim}$ in P.

The metric $G_{\lim}$ induces a metric $i^*G_{\lim}$ on $\mathcal{A}_{\lim} \cap \breve{P}$ that could be of interest to the metric approach to maximal degeneration of Calabi-Yau manifolds [KS1, KS2, Gr, GS1, GS2], in the case when P is reflexive. Even though the induced metric on the hypersurface will not in general be Ricci flat, it is not inconceivable that by carefully choosing the available parameters one might obtain the desired asymptotic behaviour.

In this respect, it is worth pointing out that a more general framework can be considered by taking a family of symplectic potentials of the form

$$g_s = g_P + \psi_s,$$

where the functions ψ_s are smooth and strictly convex in P for all $s > 0$, and where $\frac{1}{s}\psi_s \to \psi_{\text{lim}}$ as $s \to \infty$, in the C^2-norm in $C^\infty(P)$, where ψ_{lim} is strictly convex. Then, ψ_{lim} will be a potential for the limit metric. We noted already that the results in this work remain valid for this more general family of deformations.

In a different context, in [Pa1, Pa2] Parker considers degenerating families of almost complex structures in an extension of the smooth category, constructed using symplectic field theory, to study holomorphic curve invariants. The typical behaviour of his families of almost complex structures, depicted for the moment polytope in the case of $\mathbb{P}^2$ in the introduction of [Pa1], is, in the case of toric manifolds, remarkably reproduced in our approach. In fact, in the notation of sections 2 and 4,

$$\sum_{l,k=1}^{n} (G_0)_{jk}(G_0 + s\text{Hess}\,\psi)^{-1}_{kl}\frac{\partial}{\partial y_l} = \frac{\partial}{\partial y^s_j},$$

where $J_s(\frac{\partial}{\partial\theta_j}) = \frac{\partial}{\partial y^s_j}$. Therefore, in interior regions of P, where as $s \to \infty$ the term with $\text{Hess}\,\psi$ dominates, we have that the coordinates y^s appear stretched relative to the coordiantes y by an y-dependent transformation that scales with s. On the other hand, as we approach a face F of P where some coordinates l_j vanish, the derivatives of g_P with respect to these l_j's will dominate, so that the corresponding $y^s_{l_j}$'s do not scale with respect to the y_{l_j}'s. In the directions parallel to F, however, we still have have the term with $\text{Hess}\,\psi$ dominating and for these directions the scaling with s will occur. This is exactly the qualitative behaviour described in [Pa1]. We note, however, that in our approach this behaviour is implemented by deforming integrable toric complex structures.

Acknowledgements: We wish to thank Miguel Abreu for many useful conversations and for suggesting possible applications to tropical geometry. This work is partially supported by the Center for Mathematical Analysis, Geometry and Dynamical Systems, IST, by Fundação para a Ciência e a Tecnologia through the Program POCI 2010/FEDER and by the projects POCI/MAT/58549/2004 and PPCDT/MAT/58549/2004. TB is also supported by the fellowship SFRH/BD/ 22479/2005 of Fundação para a Ciência e a Tecnologia.

References

[Ab1] M.Abreu, “Kähler geometry of toric varieties and extremal metrics”, Internat. J. Math., **9** (1998), 641–651.

[Ab2] M.Abreu, “Kähler geometry of toric manifolds in symplectic coordinates”, in ”Symplectic and Contact Topology: Interactions and Perspectives” (eds. Y.Eliashberg, B.Khesin and F.Lalonde), Fields Institute Communications 35, Amer. Math. Soc., 2003.

[An] J.E.Andersen, “Jones-Witten theory and the Thurston boundary of Teichmüller spaces”, University of Oxford D. Phil. thesis, 1992.

[AdPW] S.Axelrod, S.D.Pietra and E.Witten, “Geometric quantization of Chern-Simons gauge theory”, J. Diff. Geom. **33** (1991) 787.

[BMN] T.Baier, J.Mourão and J.P.Nunes, “The BKS pairing on Abelian varieties”, in preparation.

[Bou] N. Bourbaki, “Espaces vectoriels topologiques.” Act. Sci. Ind., no. 1229. Hermann & Cie, Paris, 1955.

[BGU] D.Burns, V.Guillemin and A.Uribe, “The spectral density function of a toric variety”, arXiv:0706.3039.

[Co] D.Cox, “Minicourse on toric varieties”, University of Buenos Aires, 2001.

[Da] V.I.Danilov, “The geometry of toric varieties”, Russ. Math. Surveys **33** (1978) 97–154.

[dS] A.C. da Silva, “Symplectic toric manifolds”, in “Symplectic Geometry of Integrable Hamiltonian Systems”, Birkhäuser (Springer), 2003.

[De] T.Delzant, “Hamiltoniens periodiques et image convex de l’application moment”, Bull. Soc. Math. France, **116** (1988) 315–339.

[Do] S.Donaldson, “Scalar curvature and stability of toric varieties”, J. Diff. Geom. **62** (2002), 289–349.

[EKL] M.Einsiedler, M.Kapranov and D.Lind, “Non-Archimedean amoebas and tropical varieties”, J. Reine Angew. Math. **601** (2006) 139–157.

[FMMN] C.Florentino, P.Matias, J.Mourão and J.P.Nunes, “Geometric quantization, complex structures and the coherent state transform”, J. Funct. Anal. **221** (2005) 303–322; “On the BKS pairing for Kähler quantizations of the cotangent bundle of a Lie group”, J. Funct. Anal. **234** (2006) 180–198.

[FMN] C.Florentino, J.Mourão and J.P.Nunes, “Coherent state transforms and abelian varieties”, J. Funct. Anal. **192** (2002) 410–424; “Coherent state transforms and vector bundles on elliptic curves”, J. Funct. Anal. **204** (2003) 355–398.

[FPT] M.Forsberg, M.Passare and A.Tsikh, “Laurent determinants and arrangements of hyperplane amoebas”, Adv. Math. **151** (2000) 45–70.

[GKZ] I.M.Gelfand, M.M.Kapranov, A.V.Zelevinsky, “Discriminants, resultants and multidimensional determinants”, Birkhäuser, 1994.

[Gr] M.Gross, “The Strominger-Yau-Zaslow conjecture: from torus fibrations to degenerations”, arXiv:0802.3407.

[GS1] M.Gross and B.Siebert, “Mirror symmetry via logarithmic degeneration data. I.”, J. Diff. Geom. **72** (2006) 169–338.

[GS2] M.Gross and B.Siebert, “Mirror symmetry via logarithmic degeneration data II”, arXiv:0709.2290.

[Gui] V.Guillemin, “Kähler structures on toric varieties”, J. Diff. Geom. **40** (1994), 285–309.

[GGK] V.Guillemin, V.Ginzburg, and Y.Karshon, “Moment maps, cobordisms, and Hamiltonian group actions”, Mathematical Surveys and Monographs, 98, Amer. Math. Soc., Providence, RI, 2002.

[Gun] R.C.Gunning, “Lectures on Riemann surfaces”, Princeton University Press, 1966.

[Hal] B.C.Hall, “Geometric quantization and the generalized Segal-Bargmann transform for Lie groups of compact type”, Comm. Math. Phys. **226** (2002) 233–268.

[Ham1] M.Hamilton, “The quantization of a toric manifold is given by the integer lattice points in the moment polytope”, arXiv:0708.2710.

[Ham2] M.Hamilton, “Locally toric manifolds and singular Bohr-Sommerfeld leaves”, arXiv:0709.4058.

[Hi] N.J.Hitchin, “Flat connections and geometric quantization”, Comm. Math. Phys. **131** (1990) 347–380.

[KY] J.Kajiwara, M.Yoshida, “Note on Cauchy-Riemann equation”, Mem. Fac. Sci. Kyushu Univ. Ser. A **22** (1968) 18–22.

[Ki] A.A.Kirillov, “Geometric quantization”, in: Encyclopaedia of Mathematical Sciences, vol. 4 Dynamical systems, Springer-Verlag, 1990, 137–172.

[KW] W.Kirwin and S.Wu, “Geometric Quantization, Parallel Transport and the Fourier Transform”, Comm. Math. Phys. **266** (2006) 577-594.

[KS1] M.Kontsevich and Y.Soibelman, “Homological mirror symmetry and torus fibrations”, in “Symplectic geometry and mirror symmetry (Seoul, 2000)”, 203–263, World Sci. Publ., River Edge, NJ, 2001.

[KS2] M.Kontsevich and Y.Soibelman, “Affine structures and non-Archimedean analytic spaces”, in “The unity of mathematics”, 321–385, Progr. Math., 244, Birkhäuser Boston, Boston, MA, 2006.

[MSY] D.Martelli, J.Sparks and S.-T.Yau, “The geometric dual of a-maximisation for toric Sasaki-Einstein manifolds”, Comm. Math. Phys. **268** (2006), 39–65.

[Mi] G.Mikhalkin, "Decomposition into pairs-of-pants for complex algebraic hypersurfaces", Topol. **43** (2004), 1035–1065.

[Mu] D.Mumford, "Tata lectures on theta", Birkhäuser, 1984.

[Od] T.Oda, "Convex bodies in algebraic geometry: an introduction to toric varieties", Springer-Verlag, 1988.

[Pa1] B.Parker, "Exploded fibrations", Proceedings of the 13^{th} Gökova Geometry-Topology Conference, 1–39, arXiv:0705.2408.

[Pa2] B.Parker, "Holomorphic curves in exploded torus fibrations: compactness", arXiv:0706.3917.

[R] H.Rullgård, "Polynomial amoebas and convexity", Stockolm Univ. preprint, 2001.

[SeD] R.Sena-Dias, "Spectral measures on toric varieties and the asymptotic expansion of Tian-Yau-Zelditch", arXiv:0803.0298.

[Sn] J.Śniatycki, "Geometric quantization and quantum mechanics", Applied Math. Sciences, **30**, Springer-Verlag, 1980.

[Sn2] J.Śniatycki, "On cohomology groups appearing in geometric quantization", in: Lect. Notes in Math. **570**, Springer-Verlag, 1975.

[SYZ] A.Strominger, S.-T.Yau, E.Zaslow, "Mirror Symmetry is T-duality", Nucl. Phys. **B479** (1996), 243-259.

[Tr] F.Treves "Topological Vector Spaces, Distributions and Kernels", Academic Press, New York, 1967.

[Ty] A.Tyurin, "Quantization, field theory and theta functions", CRM Monographs, Vol.21, Amer. Math. Soc., 2003.

[Wo] N.M.J.Woodhouse, "Geometric quantization", Second Edition, Clarendon Press, Oxford, 1991.

E-mail address: `tbaier, cfloren, jmourao, jpnunes@math.ist.utl.pt`

DEPARTMENT OF MATHEMATICS, INSTITUTO SUPERIOR TÉCNICO, AV. ROVISCO PAIS, 1049-001 LISBOA, PORTUGAL